

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 April 2004 (22.04.2004)

PCT

(10) International Publication Number  
WO 2004/032861 A2

(51) International Patent Classification<sup>7</sup>: A61K  
(21) International Application Number:  
PCT/US2003/031960  
(22) International Filing Date: 9 October 2003 (09.10.2003)  
(25) Filing Language: English  
(26) Publication Language: English  
(30) Priority Data:  
60/417,935 11 October 2002 (11.10.2002) US

(71) Applicant (for all designated States except US): BRISTOL-MYERS SQUIBB COMPANY [US/US]; P. O. Box 4000, Route 206 and Provinceline Road, Princeton, NJ 08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DHAR, T., G., Murali [US/US]; 247 Crittenden Drive, Newtown, PA 18940 (US). POTIN, Dominique [FR/FR]; 14. Chemin Vert, F-78680 Epone (FR). MAILLET, Magali, Jeanine, Blandine [FR/FR]; 80 avenue de Fouilleuse, F-92150 Suresnes (FR). LAUNAY, Michele [FR/FR]; 11 rue Diderot, F-92500 Rueil Malmaison (FR). NICOLAI, Eric, Antoine [FR/FR]; 16 rue Alexis Bouvier, F-92500 Rueil Malmaison (FR). IWANOVICZ, Edwin, J. [US/US]; 4715 Dunham Court, San Diego, CA 92130 (US).

(74) Agents: DUNCAN, Laurelee et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


#### Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: HEXAHYDRO-BENZIMIDAZOLONE COMPOUNDS USEFUL AS ANTI-INFLAMMATORY AGENTS



(I)

(57) Abstract: The present invention is directed to compounds having the formula (I) useful in treating inflammatory and immune diseases, in which K is O or S; Q is -C(=O)- or optionally substituted C<sub>1-4</sub>alkylene; Ar is optionally-substituted aryl or heteroaryl; J<sub>1</sub>, J<sub>2</sub>, J<sub>3</sub>, and Y are selected so that ring A is a five-to-six membered optionally-substituted cycloalkenyl or heterocyclo ring having 0 to 2 nitrogen heteroatoms; Z is N or C(R<sub>9</sub>); and R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>9</sub> are as defined in the specification.

WO 2004/032861 A2



**Published:**

- without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## HEXAHYDRO-BENZIMIDAZOLONE COMPOUNDS USEFUL AS ANTI-INFLAMMATORY AGENTS

This application claims benefit of priority from U.S. Provisional Application  
5 Serial No. 60/417,935 filed October 11, 2002, the entirety of which is incorporated  
herein by reference.

### Field of the Invention

The present invention relates to hexahydro-benzimidazolone compounds,  
pharmaceutical compositions containing them, and methods of using such compounds  
10 in treating inflammatory or immune disease.

### Background of the Invention

Cells adhere to other cells and to substrates through specific, regulated  
processes that are critical to various biological functions. The proper functioning of  
the immune system is dependent upon adhesive interactions and cell migration. A key  
15 event in an immune response involves the migration of leukocytes to a disease site.  
During an inflammatory response, leukocytes are recruited to the site of injury and  
extravasated by a series of cellular interactions involving cell-cell and cell-substrate  
adhesion.

One family of molecules that serves an important adhesive function is  
20 integrins. Integrins are expressed on cell surfaces and function in cell-cell and cell-  
substrate adhesion. Integrins are alpha-beta heterodimers: each integrin has an alpha  
( $\alpha$ ) subunit non-covalently bound to a beta ( $\beta$ ) subunit. When activated, integrins bind  
to extracellular ligands and induce adhesion (the expression of integrins on a cell  
surface alone is inadequate for adhesion - they must be activated to become adhesive).  
25 The integrin activation state is transient, such that there is a rapid flux between  
adhesive and non-adhesive states which is important for cell movement, *e.g.*, a cell is  
endowed with the ability to rapidly adhere to various cell surfaces and matrices and to  
migrate among cells and tissue.

There are four known integrins having a  $\beta_2$  or CD18 subunit which comprise  
30 the CD11/CD18 integrin sub-family, namely, Lymphocyte Function-associated  
Antigen 1 (LFA-1) (CD11a/CD18 or  $\alpha_L\beta_2$ ); Macrophage Antigen 1 (Mac-1)

(CD11b/CD18 or  $\alpha_M\beta_2$ ); p150,95 (CD11c/CD18 or  $\alpha_X\beta_2$ ); and  $\alpha_D\beta_2$ . The CD11/CD18 family of integrins is also referred to as Leukointegrins as they are expressed on the surface of various leukocyte cells, and they mediate a number of inflammation-related cellular interactions. See Diamond *et al.*, "The Dynamic  
5 Regulation of Integrin Adhesiveness," *Current Biology*, Vol. 4 (1994) at pp. 506-532.

Ligands to LFA-1 and Mac-1 comprise the intercellular adhesion molecule (ICAM) ICAM-1. The primary CD11/CD18 integrin is LFA-1, which also binds with ICAM-2 and ICAM-3. ICAMs are found on endothelium cells, leukocytes, and other cell types, and their interaction with CD11/CD18 integrins is critical to immune  
10 system function. The interaction between the CD18 integrins, particularly LFA-1, and ICAMs mediates antigen presentation, T-cell proliferation, and adhesion between the endothelium and activated leukocytes which is necessary for leukocytes to migrate from the circulatory system into tissue. A condition termed "Leukocyte Adhesion Deficiency" has been identified in patients having a deficiency in CD18 integrins.  
15 These patients are unable to mount a normal inflammatory or immune response; they suffer from disorders such as recurrent infections, poor wound healing, granulocytosis, progressive periodontitis, and umbilical cord separation. See Anderson *et al.*, "Leukocyte LFA-1, OKMI, p150,95 Deficiency Syndrome: Functional and Biosynthesis Studies of Three Kindreds," *Fed. Proc.*, Vol. 44 (1985), at pp. 2671-  
20 2677.

While sufficient levels of CD18 integrins interacting with ICAMs are needed to mount a normal immune response, significant cellular and tissue injury can result in chronic inflammatory states where there is an inappropriate influx of leukocytes to the disease site. Continuous recruitment of leukocytes from blood vessels into inflamed  
25 tissue, as in chronic inflammatory states, can perpetuate tissue injury and lead to excessive fibrous repair and autoimmune disease. Thus, inhibition of the interaction between LFA-1 and/or Mac-1 and their ICAMs can be advantageous in treating inflammatory or immune disease. For example, monoclonal antibody blockade of either ICAM or LFA-1 has been shown to prevent the migration of leukocytes into  
30 tissue and the subsequent development of inflammatory disease in animal models of rheumatoid arthritis, inflammatory bowel disease, and pulmonary inflammation (*e.g.*, asthma). Knockout mice deficient in ICAMs have reduced susceptibility to induced

arthritis, ischemia injury, impaired lung inflammatory responses, and increased tolerance to transplantations (e.g. heart grafts). See Anderson, *supra*. Antibodies blocking the ICAM-LFA-1 interaction reportedly suppress cardiac allograft rejection and islet cell xenograft rejection in animal models. See Gorski, "The Role of Cell  
5 *Adhesion Molecules in Immunopathology*," *Immunology Today*, Vol. 15 (1994), at pp. 251-255.

Compounds inhibiting CD18 integrins, ICAMs, and/or the LFA-1:ICAM interaction could potentially demonstrate a wide range of utilities in treating inflammatory or immune diseases. Blocking LFA-1 reportedly inhibits the influx of  
10 leukocytes in almost every system, including the skin, peritoneum, synovium, lung, kidney, and heart, and blocking ICAM-1 would be expected to have similar effects. Also, present therapies for many inflammatory and immune diseases have drawbacks. For example, current treatments for asthma include  $\beta_2$ -agonists, inhaled corticosteroids, and LTD<sub>4</sub> antagonists. However,  $\beta_2$ -agonists have limited efficacy  
15 and inhaled corticosteroids raise safety concerns. To treat psoriasis, current therapies include PUVA, methotrexate, cyclosporin A, and topical treatments. The first three of these therapies raise toxicity issues over long-term (e.g., 6-9 month) use, whereas topical treatments have limited efficacy. Additionally, these treatments typically are applied only in response to flares and not as a prophylaxis measure.

20 Compounds that reportedly inhibit LFA-1/ICAM for use as anti-inflammatory agents include thiadiazole-based compounds (see Intern. Pub. No. WO 99/20,618, "Thiadiazole Amides Useful as Anti-Inflammatory Agents" filed by Pharmacia & Upjohn Co.; and WO 99/20,617, also to Pharmacia and Upjohn). Small molecules that reportedly are antagonists to the binding of ICAMs with CD18 integrins include  
25 various benzylamines and 2-bromobenzoyltryptophan compounds (see Intern. Pub. No. WO99/49,856), and 1-(3,5 dichlorophenyl) imidazolidines (see Intern. Pub. No. WO98/39303, "Small Molecules Useful in the Treatment of Inflammatory Disease," filed by Boehringer Ingelheim Pharmaceuticals, Inc. See also Boehringer patent applications WO 01/07052, WO 01/07048, WO 01/07044, WO 01/06984, and WO  
30 01/07440). Hydantoin compounds are disclosed in Intern. Pub. No's WO 00/59880, WO 00/39081, WO 02/02522, WO 02/02539 (all to Abbott Laboratories). LFA-1 antagonist compounds are also claimed in WO 02/059114 (to Genentech), WO


02/42294 (to Celltech), WO 01/51508 (to Science and Technology corporation), WO 00/21920 and WO 01/58853 (both to Hoffmann-LaRoche), and WO 99/11258, WO 00/48989 and WO 02/28832 (all to Novartis). Hydantoin compounds are disclosed in WO 01/30781 A2 (published May 3, 2001) to Tanabe Seiyaku Co. Ltd, "*Inhibitors of*  
5  *$\alpha_L\beta_2$  Mediated Cell Adhesion*". See also, WO 02/44181 (published June 6, 2002), "*Hydantoin Compounds Useful as Anti-Inflammatory Agents*", and Attny Docket No. QA263 filed October 2, 2003, to the present assignee and having common inventors herewith.

As may be appreciated, those in the field of pharmaceutical research continue  
10 to develop new compounds and compositions for treating inflammatory and immune disease such as inhibitors of Leukointegrins and/or ICAMs. Particularly in the area of immune response, many individuals respond differently to different drugs. Thus, there is an interest in providing consumers not only with pharmaceutical compositions demonstrating increased effectiveness and reduced side-effects but also different  
15 structures or mechanisms of action to provide consumers with a choice of options. The instant invention is directed to aryl substituted hexahydro-benzimidazolone compounds that are effective as antagonists of Leukointegrins and/or ICAMs. Each of the patents, patent applications and publications referenced above and hereinafter is incorporated herein by reference.

20

### Summary of the Invention

The present invention provides compounds useful in treating inflammatory or immune disease having the formula (I):



25 its enantiomers, diastereomers, and pharmaceutically-acceptable salts, hydrates, and prodrugs thereof, in which:

K is O or S;

Q is a bond,  $-C(=O)-$  or branched or straight chain  $C_{1-4}$ alkylene optionally substituted with one to two  $R_4$ ;

Ar is optionally-substituted aryl or heteroaryl;

$J_1$  is a bond,  $-N(R_5)-$ , or  $-C(R_{6a}R_{7a})-$ ;

5  $J_2$  is  $-N(R_5)-$  or  $-C(R_{6b}R_{7b})-$ ;

$J_3$  is  $-N(R_5)-$  or  $-C(R_{6c}R_{7c})-$ ;

provided, however, that only one of  $J_1$ ,  $J_2$  and  $J_3$  may be  $-N(R_5)-$ , so that ring A is a five-to-six membered cycloalkyl or heterocyclo ring having from 0 to 2 heteroatoms;

10 Y is N or  $C(R_8)$ ;

Z is N or  $C(R_9)$ ;

$R_1$  is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl,  $-OR_{10}$ ,  $-NR_{10}R_{11}$ ,  $-C(=O)R_{10}$ ,  $-CO_2R_{10}$ ,  $-C(=O)NR_{10}R_{11}$ ,  $-S(O)_pR_{11a}$ ,  $-SO_2NR_{10}R_{11}$ , cycloalkyl, heterocyclo, aryl, and heteroaryl;

15  $R_2$  and  $R_3$  are independently selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl,  $-SR_{12}$ ,  $-OR_{12}$ ,  $-NR_{12}R_{13}$ ,  $-CO_2R_{12}$ ,  $-C(=O)R_{12}$ ,  $-C(=O)NR_{12}R_{13}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl;

20  $R_4$  is selected from OH,  $O(C_{1-4}alkyl)$ , halogen, cyano,  $CF_3$ ,  $OCF_3$ ,  $NH_2$ ,  $NH(C_{1-4}alkyl)$ , and  $N(C_{1-4}alkyl)_2$ ;

$R_5$  is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cyano,  $-OR_{14}$ ,  $-NR_{14}R_{15}$ ,  $-C(=O)R_{14}$ ,  $-CO_2R_{14}$ ,  $-C(=O)NR_{14}R_{15}$ ,  $-S(O)_pR_{15a}$ ,  $-SO_2NR_{14}R_{15}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl; or when  $R_5$  is joined to atom  $J_1$ ,  $J_2$  or  $J_3$ ,  $R_5$  may be taken together with one of  $R_{6a}$ ,  $R_{6b}$  or  $R_{6c}$  attached to an adjacent atom of ring A to form a fused heterocyclo or heteroaryl ring; or when  $R_5$  is joined to atom  $J_3$ ,  $R_5$  may be taken together with  $R_8$  to form a fused heterocyclo ring;

25

$R_{6a}$ ,  $R_{6b}$ ,  $R_{6c}$ ,  $R_{7a}$ ,  $R_{7b}$ ,  $R_{7c}$  and  $R_8$  are independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, nitro, cyano,  $-SR_{16}$ , -

OR<sub>16</sub>, -NR<sub>16</sub>R<sub>17</sub>, -C(=O)R<sub>16</sub>, -CO<sub>2</sub>R<sub>16</sub>, -C(=O)NR<sub>16</sub>R<sub>17</sub>, -NR<sub>16</sub>C(=O)R<sub>17</sub>, -  
NR<sub>16</sub>C(=O)OR<sub>17</sub>, -S(O)<sub>q</sub>R<sub>17a</sub>, -NR<sub>16</sub>SO<sub>2</sub>R<sub>17a</sub>, -SO<sub>2</sub>NR<sub>16</sub>R<sub>17</sub>, aryl, heterocyclo,  
cycloalkyl, and heteroaryl; or R<sub>6a</sub> with R<sub>7a</sub>, or R<sub>6b</sub> with R<sub>7b</sub>, or R<sub>6c</sub> with R<sub>7c</sub> are  
taken together to form a keto group (=O) or a spiro cycloalkyl or heterocyclo  
5 ring; or R<sub>6b</sub> taken together with either R<sub>6a</sub> or R<sub>6c</sub> may form a fused benzo,  
cycloalkyl, heterocyclo, or heteroaryl ring; or R<sub>6c</sub> taken together with R<sub>8</sub> may  
form a fused cycloalkyl or heterocyclo;

R<sub>9</sub> is selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl,  
substituted alkenyl, -SR<sub>18</sub>, -OR<sub>18</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>18</sub>, -C(=O)R<sub>18</sub>, -  
10 C(=O)NR<sub>18</sub>R<sub>19</sub>, aryl, heterocyclo, cycloalkyl, and heteroaryl;

R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>, and R<sub>19</sub> (i) are selected independently of  
each other from hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl,  
and heterocyclo; or (ii) any two of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>,  
and R<sub>19</sub> when attached to the same nitrogen atom (as in NR<sub>10</sub>R<sub>11</sub>, NR<sub>12</sub>R<sub>13</sub>,  
15 NR<sub>14</sub>R<sub>15</sub>, NR<sub>16</sub>R<sub>17</sub>, or NR<sub>18</sub>R<sub>19</sub>) may be taken together to form a heteroaryl or  
heterocyclo ring, with the remainder of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>,  
R<sub>18</sub>, and R<sub>19</sub> being selected independently from hydrogen, alkyl, substituted  
alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo;

R<sub>11a</sub>, R<sub>15a</sub>, and R<sub>17a</sub> are independently selected from alkyl, substituted alkyl,  
20 cycloalkyl, aryl, heteroaryl, and heterocyclo;

*p* is 1, 2, or 3; and

*q* is 1, 2, or 3.

The present invention is also directed to pharmaceutical compositions useful  
25 in treating immune or inflammatory diseases comprising compounds of formula (I), or  
pharmaceutically-acceptable salts thereof, and pharmaceutically-acceptable carriers or  
diluent. The invention further relates to methods of treating immune or inflammatory  
diseases comprising administering to a patient in need of such treatment a  
therapeutically-effective amount of a compound according to formula (I).



### Detailed Description of the Invention

The following are definitions of terms used in this specification and appended claims. The initial definition provided for a group or term herein applies to that group or term throughout the specification and claims, individually or as part of another group, unless otherwise indicated.

The term "alkyl" refers to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms. Lower alkyl groups, that is, alkyl groups of 1 to 4 carbon atoms, are most preferred. When numbers appear in a subscript after the symbol "C", the subscript defines with more specificity the number of carbon atoms that a particular group may contain. For example, "C<sub>1-6</sub>alkyl" refers to straight and branched chain alkyl groups with one to six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, and so forth. The subscript "0" refers to a bond. Thus, the term hydroxy(C<sub>0-2</sub>)alkyl or (C<sub>0-2</sub>)hydroxyalkyl includes hydroxy, hydroxymethyl and hydroxyethyl.

The term "substituted alkyl" refers to an alkyl group as defined above having one, two, or three substituents selected from the group consisting of halo (*e.g.*, trifluoromethyl), alkenyl, substituted alkenyl, alkynyl, nitro, cyano, oxo (=O), OR<sub>a</sub>, SR<sub>a</sub>, (=S), -NR<sub>a</sub>R<sub>b</sub>, -N(alkyl)<sub>3</sub><sup>+</sup>, -NR<sub>a</sub>SO<sub>2</sub>, -NR<sub>a</sub>SO<sub>2</sub>R<sub>c</sub>, -SO<sub>2</sub>R<sub>c</sub>, -SO<sub>2</sub>NR<sub>a</sub>R<sub>b</sub>, -SO<sub>2</sub>NR<sub>a</sub>C(=O)R<sub>b</sub>, SO<sub>3</sub>H, -PO(OH)<sub>2</sub>, -C(=O)R<sub>a</sub>, -CO<sub>2</sub>R<sub>a</sub>, -C(=O)NR<sub>a</sub>R<sub>b</sub>, -C(=O)(C<sub>1-4</sub>alkylene)NR<sub>a</sub>R<sub>b</sub>, -C(=O)NR<sub>a</sub>(SO<sub>2</sub>)R<sub>b</sub>, -CO<sub>2</sub>(C<sub>1-4</sub>alkylene)NR<sub>a</sub>R<sub>b</sub>, -NR<sub>a</sub>C(=O)R<sub>b</sub>, -NR<sub>a</sub>CO<sub>2</sub>R<sub>b</sub>, -NR<sub>a</sub>(C<sub>1-4</sub>alkylene)CO<sub>2</sub>R<sub>b</sub>, =N-OH, =N-O-alkyl, aryl, cycloalkyl, heterocyclo, and/or heteroaryl, wherein R<sub>a</sub> and R<sub>b</sub> are selected from hydrogen, alkyl, alkenyl, CO<sub>2</sub>H, CO<sub>2</sub>(alkyl), C<sub>3-7</sub>cycloalkyl, phenyl, benzyl, phenylethyl, naphthyl, a four to seven membered heterocyclo, or a five to six membered heteroaryl, or when attached to the same nitrogen atom may join to form a heterocyclo or heteroaryl, and R<sub>c</sub> is selected from same groups as R<sub>a</sub> and R<sub>b</sub> but is not hydrogen. Each group R<sub>a</sub> and R<sub>b</sub> when other than hydrogen, and each R<sub>c</sub> group optionally has up to three further substituents attached at any available carbon or nitrogen atom of R<sub>a</sub>, R<sub>b</sub>, and/or R<sub>c</sub>, said substituent(s) being selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>2-6</sub>)alkenyl, hydroxy, halogen, cyano, nitro, CF<sub>3</sub>, O(C<sub>1-6</sub>alkyl), OCF<sub>3</sub>, C(=O)H, C(=O)(C<sub>1-6</sub>alkyl), CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-6</sub>alkyl), NHCO<sub>2</sub>(C<sub>1-6</sub>alkyl), -S(C<sub>1-6</sub>alkyl), -NH<sub>2</sub>, NH(C<sub>1-6</sub>alkyl), N(C<sub>1-6</sub>alkyl)<sub>2</sub>, N(CH<sub>3</sub>)<sub>3</sub><sup>+</sup>, SO<sub>2</sub>(C<sub>1-6</sub>alkyl), C(=O)(C<sub>1-4</sub>alkylene)NH<sub>2</sub>,

C(=O)(C<sub>1-4</sub>alkylene)NH(alkyl), C(=O)(C<sub>1-4</sub>alkylene)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, C<sub>3-7</sub>cycloalkyl, phenyl, benzyl, phenylethyl, phenyloxy, benzyloxy, naphthyl, a four to seven membered heterocyclo, or a five to six membered heteroaryl. When a substituted alkyl is substituted with an aryl, heterocyclo, cycloalkyl, or heteroaryl group, said ringed systems are as defined below and thus may have zero, one, two, or three substituents, also as defined below.

One skilled in the field will understand that, when the designation "CO<sub>2</sub>" is used herein, this is intended to refer to the group  $\text{---}\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{---O---}$ .

When the term "alkyl" is used together with another group, such as in "arylalkyl", this conjunction defines with more specificity at least one of the substituents that the substituted alkyl will contain. For example, "arylalkyl" refers to a substituted alkyl group as defined above where at least one of the substituents is an aryl, such as benzyl. Thus, the term aryl(C<sub>0-4</sub>)alkyl includes a substituted lower alkyl having at least one aryl substituent and also includes an aryl directly bonded to another group, *i.e.*, aryl(C<sub>0</sub>)alkyl.

The term "alkenyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms and at least one double bond. Alkenyl groups of 2 to 6 carbon atoms and having one double bond are most preferred.

The term "alkynyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms and at least one triple bond. Alkynyl groups of 2 to 6 carbon atoms and having one triple bond are most preferred.

The term "alkylene" refers to bivalent straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms, *e.g.*, {-CH<sub>2</sub>-}<sub>n</sub>, wherein *n* is 1 to 12, preferably 1-8. Lower alkylene groups, that is, alkylene groups of 1 to 4 carbon atoms, are most preferred. The terms "alkenylene" and "alkynylene" refer to bivalent radicals of alkenyl and alkynyl groups, respectively, as defined above.

When reference is made to a substituted alkenyl, alkynyl, alkylene, alkenylene, or alkynylene group, these groups are substituted with one to three substituents as defined above for substituted alkyl groups.

The term "alkoxy" refers to an alkyl group as defined above bonded through an oxygen atom (-O-). The term "substituted alkoxy" includes the groups -O-R, wherein R is a substituted alkyl group as defined above.

The term "thioalkyl" refers to an alkyl group as defined above bonded through a sulfur atom. The term "substituted thioalkyl" includes the groups -S-R, wherein R is a substituted alkyl group as defined above.


The term "alkylamino" refers to an alkyl group as defined above bonded through a nitrogen atom. For example, the term "alkylamino" includes groups such as -NR-C<sub>1-12</sub>alkyl (where R is preferably hydrogen but may include alkyl.) "Aminoalkyl" refers to an amino group bonded through an alkyl, e.g., -(CH<sub>2</sub>)<sub>x</sub>-NH<sub>2</sub>. "Alkylaminoalkyl" refers to an alkylamino group (i.e., -NHR or NRR) bonded through an alkyl group (e.g., -(CH<sub>2</sub>)<sub>x</sub>-NHR or -(CH<sub>2</sub>)<sub>x</sub>-NRR).

When a subscript is used with reference to an alkoxy, thioalkyl, alkylamino, or aminoalkyl, the subscript refers to the number of carbon atoms that the group may contain in addition to heteroatoms. Thus, for example, monovalent C<sub>1-2</sub>alkylamino includes the groups -NH-CH<sub>3</sub>, -NH-CH<sub>2</sub>-CH<sub>3</sub>, and -N-(CH<sub>3</sub>)<sub>2</sub>.

The term "cycloalkyl" refers to fully saturated and partially unsaturated hydrocarbon rings of 3 to 9, preferably 3 to 7 carbon atoms. The term "cycloalkyl" includes such rings having zero, one, two, or three substituents selected from the group consisting of halogen, trifluoromethyl, trifluoromethoxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, nitro, cyano, oxo (=O), OR<sub>a</sub>, SR<sub>a</sub>, (=S), -NR<sub>a</sub>R<sub>b</sub>, -N(alkyl)<sub>3</sub><sup>+</sup>, -NR<sub>a</sub>SO<sub>2</sub>, -NR<sub>a</sub>SO<sub>2</sub>R<sub>c</sub>, -SO<sub>2</sub>R<sub>c</sub>, -SO<sub>2</sub>NR<sub>a</sub>R<sub>b</sub>, -SO<sub>2</sub>NR<sub>a</sub>C(=O)R<sub>b</sub>, SO<sub>3</sub>H, -PO(OH)<sub>2</sub>, -C(=O)R<sub>a</sub>, -CO<sub>2</sub>R<sub>a</sub>, -C(=O)NR<sub>a</sub>R<sub>b</sub>, -C(=O)(C<sub>1-4</sub>alkylene)NR<sub>a</sub>R<sub>b</sub>, -C(=O)NR<sub>a</sub>(SO<sub>2</sub>)R<sub>b</sub>, -CO<sub>2</sub>(C<sub>1-4</sub>alkylene)NR<sub>a</sub>R<sub>b</sub>, -NR<sub>a</sub>C(=O)R<sub>b</sub>, -NR<sub>a</sub>CO<sub>2</sub>R<sub>b</sub>, -NR<sub>a</sub>(C<sub>1-4</sub>alkylene)CO<sub>2</sub>R<sub>b</sub>, =N-OH, =N-O-alkyl, aryl, cycloalkyl, heterocyclo, and/or heteroaryl, wherein R<sub>a</sub>, R<sub>b</sub> and R<sub>c</sub> are as defined above for substituted alkyl groups, and are also in turn optionally substituted as recited above in the definition for substituted alkyl groups. The term "cycloalkyl" also includes such rings having a second ring fused thereto (e.g., including benzo, heterocyclo, or heteroaryl rings) or having a carbon-carbon bridge of 3 to 4 carbon atoms. When a cycloalkyl is substituted with a further ring (or has a second ring fused thereto), said ring in turn is

optionally substituted with one to two of (C<sub>1-4</sub>)alkyl, (C<sub>2-4</sub>)alkenyl, halogen, hydroxy, cyano, nitro, CF<sub>3</sub>, O(C<sub>1-4</sub>alkyl), OCF<sub>3</sub>, C(=O)H, C(=O)(C<sub>1-4</sub>alkyl), CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-4</sub>alkyl), NHCO<sub>2</sub>(C<sub>1-4</sub>alkyl), -S(C<sub>1-4</sub>alkyl), -NH<sub>2</sub>, NH(C<sub>1-4</sub>alkyl), N(C<sub>1-4</sub>alkyl)<sub>2</sub>, N(C<sub>1-4</sub>alkyl)<sub>3</sub><sup>+</sup>, SO<sub>2</sub>(C<sub>1-4</sub>alkyl), C(=O)(C<sub>1-4</sub>alkylene)NH<sub>2</sub>, C(=O)(C<sub>1-4</sub>alkylene)NH(alkyl),  
 5 and/or C(=O)(C<sub>1-4</sub>alkylene)N(C<sub>1-4</sub>alkyl)<sub>2</sub>.

Accordingly, in compounds of formula (I), the term "cycloalkyl" includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc., as well as the following ring systems,



The term "halo" or "halogen" refers to chloro, bromo, fluoro and iodo.

15 The term "haloalkyl" means a substituted alkyl having one or more halo substituents. For example, "haloalkyl" includes mono, bi, and trifluoromethyl.

The term "haloalkoxy" means an alkoxy group having one or more halo substituents. For example, "haloalkoxy" includes OCF<sub>3</sub>.


20 The term "aryl" refers to phenyl, biphenyl, 1-naphthyl and 2-naphthyl. The term "aryl" includes such rings having zero, one, two or three substituents selected from the group consisting of halogen, trifluoromethyl, trifluoromethoxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, nitro, cyano, OR<sub>a</sub>, SR<sub>a</sub>, (=S), -NR<sub>a</sub>R<sub>b</sub>, -N(alkyl)<sub>3</sub><sup>+</sup>, -NR<sub>a</sub>SO<sub>2</sub>, -NR<sub>a</sub>SO<sub>2</sub>R<sub>c</sub>, -SO<sub>2</sub>R<sub>c</sub> -SO<sub>2</sub>NR<sub>a</sub>R<sub>b</sub>, -SO<sub>2</sub>NR<sub>a</sub>C(=O)R<sub>b</sub>, SO<sub>3</sub>H, -PO(OH)<sub>2</sub>, -C(=O)R<sub>a</sub>, -CO<sub>2</sub>R<sub>a</sub>, -C(=O)NR<sub>a</sub>R<sub>b</sub>, -C(=O)(C<sub>1-4</sub>alkylene)NR<sub>a</sub>R<sub>b</sub>, -

$C(=O)NR_a(SO_2)R_b$ ,  $-CO_2(C_{1-4}alkylene)NR_aR_b$ ,  $-NR_aC(=O)R_b$ ,  $-NR_aCO_2R_b$ ,  $-NR_a(C_{1-4}alkylene)CO_2R_b$ , aryl, cycloalkyl, heterocyclo, and/or heteroaryl, wherein  $R_a$ ,  $R_b$  and  $R_c$  are as defined above for substituted alkyl groups, and are also in turn optionally substituted as recited above. Additionally, two substituents attached to an aryl,

5 particularly a phenyl group, may join to form a further ring such as a fused or spiro-ring, *e.g.*, cyclopentyl or cyclohexyl, or fused heterocyclo or heteroaryl. When an aryl is substituted with a further ring (or has a second ring fused thereto), said ring in turn is optionally substituted with one to two of  $(C_{1-4})alkyl$ ,  $(C_{2-4})alkenyl$ , halogen, hydroxy, cyano, nitro,  $CF_3$ ,  $O(C_{1-4}alkyl)$ ,  $OCF_3$ ,  $C(=O)H$ ,  $C(=O)(C_{1-4}alkyl)$ ,  $CO_2H$ ,

10  $CO_2(C_{1-4}alkyl)$ ,  $NHCO_2(C_{1-4}alkyl)$ ,  $-S(C_{1-4}alkyl)$ ,  $-NH_2$ ,  $NH(C_{1-4}alkyl)$ ,  $N(C_{1-4}alkyl)_2$ ,  $N(C_{1-4}alkyl)_3^+$ ,  $SO_2(C_{1-4}alkyl)$ ,  $C(=O)(C_{1-4}alkylene)NH_2$ ,  $C(=O)(C_{1-4}alkylene)NH(alkyl)$ , and/or  $C(=O)(C_{1-4}alkylene)N(C_{1-4}alkyl)_2$ .

Thus, examples of aryl groups include:



The terms "heterocyclo" or "heterocyclic" refers to substituted and unsubstituted non-aromatic 3 to 7 membered monocyclic groups, 7 to 11 membered

20 bicyclic groups, and 10 to 15 membered tricyclic groups, in which at least one of the rings has at least one heteroatom (O, S or N). Each ring of the heterocyclo group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less, and further provided that the ring contains at least one carbon atom.

25 The fused rings completing bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms may optionally be

quaternized. The heterocyclo group may be attached at any available nitrogen or carbon atom. The heterocyclo ring may contain zero, one, two or three substituents selected from the group consisting of halogen, trifluoromethyl, trifluoromethoxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, nitro, cyano, oxo (=O),

5  $OR_a$ ,  $SR_a$ , (=S),  $-NR_aR_b$ ,  $-N(alkyl)_3^+$ ,  $-NR_aSO_2$ ,  $-NR_aSO_2R_c$ ,  $-SO_2R_c$ ,  $-SO_2NR_aR_b$ ,  $-SO_2NR_aC(=O)R_b$ ,  $SO_3H$ ,  $-PO(OH)_2$ ,  $-C(=O)R_a$ ,  $-CO_2R_a$ ,  $-C(=O)NR_aR_b$ ,  $-C(=O)(C_{1-4}alkylene)NR_aR_b$ ,  $-C(=O)NR_a(SO_2)R_b$ ,  $-CO_2(C_{1-4}alkylene)NR_aR_b$ ,  $-NR_aC(=O)R_b$ ,  $-NR_aCO_2R_b$ ,  $-NR_a(C_{1-4}alkylene)CO_2R_b$ ,  $=N-OH$ ,  $=N-O-alkyl$ , aryl, cycloalkyl, heterocyclo, and/or heteroaryl, wherein  $R_a$ ,  $R_b$  and  $R_c$  are as defined above for


10 substituted alkyl groups, and are also in turn optionally substituted as recited above. When a heterocyclo is substituted with a further ring, said ring in turn is optionally substituted with one to two of  $(C_{1-4})alkyl$ ,  $(C_{2-4})alkenyl$ , halogen, hydroxy, cyano, nitro,  $CF_3$ ,  $O(C_{1-4}alkyl)$ ,  $OCF_3$ ,  $C(=O)H$ ,  $C(=O)(C_{1-4}alkyl)$ ,  $CO_2H$ ,  $CO_2(C_{1-4}alkyl)$ ,  $NHCO_2(C_{1-4}alkyl)$ ,  $-S(C_{1-4}alkyl)$ ,  $-NH_2$ ,  $NH(C_{1-4}alkyl)$ ,  $N(C_{1-4}alkyl)_2$ ,  $N(C_{1-4}alkyl)_3^+$ ,

15  $SO_2(C_{1-4}alkyl)$ ,  $C(=O)(C_{1-4}alkylene)NH_2$ ,  $C(=O)(C_{1-4}alkylene)NH(alkyl)$ , and/or  $C(=O)(C_{1-4}alkylene)N(C_{1-4}alkyl)_2$ .

Exemplary monocyclic groups include azetidiny, pyrrolidinyl, oxetanyl, imidazolinyl, oxazolidinyl, isoxazolinyl, thiazolidinyl, isothiazolidinyl, tetrahydrofuranyl, piperidyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidyl, 2-

20 oxopyrrolodiny, 2-oxoazepinyl, azepinyl, 4-piperidonyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl and the like. Exemplary bicyclic heterocyclo groups include quinuclidinyl.

Preferred heterocyclo groups in compounds of formula (I) include



The term "heteroaryl" refers to substituted and unsubstituted aromatic 5 or 6 membered monocyclic groups, 9 or 10 membered bicyclic groups, and 11 to 14 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heteroaryl group containing a heteroatom can

5 contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less and each ring has at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms may optionally be oxidized and the

10 nitrogen atoms may optionally be quaternized. Heteroaryl groups which are bicyclic or tricyclic must include at least one fully aromatic ring but the other fused ring or rings may be aromatic or non-aromatic. The heteroaryl group may be attached at any available nitrogen or carbon atom of any ring. The heteroaryl ring system may contain zero, one, two or three substituents selected from the group consisting of halogen,

15 trifluoromethyl, trifluoromethoxy, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, nitro, cyano,  $OR_a$ ,  $SR_a$ ,  $(=S)$ ,  $-NR_aR_b$ ,  $-N(alkyl)_3^+$ ,  $-NR_aSO_2$ ,  $-NR_aSO_2R_c$ ,  $-SO_2R_c-SO_2NR_aR_b$ ,  $-SO_2NR_aC(=O)R_b$ ,  $SO_3H$ ,  $-PO(OH)_2$ ,  $-C(=O)R_a$ ,  $-CO_2R_a$ ,  $-C(=O)NR_aR_b$ ,  $-C(=O)(C_{1-4}alkylene)NR_aR_b$ ,  $-C(=O)NR_a(SO_2)R_b$ ,  $-CO_2(C_{1-4}alkylene)NR_aR_b$ ,  $-NR_aC(=O)R_b$ ,  $-NR_aCO_2R_b$ ,  $-NR_a(C_{1-4}alkylene)CO_2R_b$ , aryl,

20 cycloalkyl, heterocyclo, and/or heteroaryl, wherein  $R_a$ ,  $R_b$  and  $R_c$  are as defined above for substituted alkyl groups, and are also in turn optionally substituted as recited above. When a heteroaryl is substituted with a further ring, said ring in turn is optionally substituted with one to two of  $(C_{1-4})alkyl$ ,  $(C_{2-4})alkenyl$ , halogen, hydroxy, cyano, nitro,  $CF_3$ ,  $O(C_{1-4}alkyl)$ ,  $OCF_3$ ,  $C(=O)H$ ,  $C(=O)(C_{1-4}alkyl)$ ,  $CO_2H$ ,  $CO_2(C_{1-4}alkyl)$ ,  $NHCO_2(C_{1-4}alkyl)$ ,  $-S(C_{1-4}alkyl)$ ,  $-NH_2$ ,  $NH(C_{1-4}alkyl)$ ,  $N(C_{1-4}alkyl)_2$ ,  $N(C_{1-4}alkyl)_3^+$ ,  $SO_2(C_{1-4}alkyl)$ ,  $C(=O)(C_{1-4}alkylene)NH_2$ ,  $C(=O)(C_{1-4}alkylene)NH(alkyl)$ , and/or  $C(=O)(C_{1-4}alkylene)N(C_{1-4}alkyl)_2$ .


Exemplary monocyclic heteroaryl groups include pyrrolyl, pyrazolyl, pyrazolinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl,

30 furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl and the like.

Exemplary bicyclic heteroaryl groups include indolyl, benzothiazolyl, benzodioxolyl, benzoxazolyl, benzothienyl, quinoliny, tetrahydroisoquinoliny, isoquinoliny, benzimidazolyl, benzopyranyl, indoliziny, benzofuranyl, chromonyl, coumariny, benzopyranyl, cinnoliny, quinoxaliny, indazolyl, pyrrolopyridyl, furopyridyl, dihydroisoindolyl, tetrahydroquinoliny and the like.

Exemplary tricyclic heteroaryl groups include carbazolyl, benzidolyl, phenanthrolliny, acridiny, phenanthridiny, xanthenyl and the like.

In compounds of formula (I), preferred heteroaryl groups include



optionally may be substituted at any available carbon or nitrogen atom.

Unless otherwise indicated, when reference is made to a specifically-named aryl (*e.g.*, phenyl), cycloalkyl (*e.g.*, cyclohexyl), heterocyclo (*e.g.*, pyrrolidiny) or heteroaryl (*e.g.*, imidazolyl), unless otherwise specifically indicated the reference is intended to include rings having 0 to 3, preferably 0-2, substituents selected from those recited above for the aryl, cycloalkyl, heterocyclo and/or heteroaryl groups, as appropriate.

The term "heteroatoms" shall include oxygen, sulfur and nitrogen.

20 The term "carbocyclic" means a saturated or unsaturated monocyclic or bicyclic ring in which all atoms of all rings are carbon. Thus, the term includes cycloalkyl and aryl rings. The carbocyclic ring may be substituted in which case the substituents are selected from those recited above for cycloalkyl and aryl groups.

25 When the term "unsaturated" is used herein to refer to a ring or group, the ring or group may be fully unsaturated or partially unsaturated.



Throughout the specification, groups and substituents thereof may be chosen by one skilled in the field to provide stable moieties and compounds and compounds useful as pharmaceutically-acceptable compounds and/or intermediate compounds useful in making pharmaceutically-acceptable compounds.

5           The compounds of formula (I) can form salts which are also within the scope of this invention. Unless otherwise indicated, reference to an inventive compound is understood to include reference to salts thereof. The term "salt(s)" denotes acidic and/or basic salts formed with inorganic and/or organic acids and bases. In addition, the term "salt(s)" may include zwitterions (inner salts), *e.g.*, when a compound of  
10   formula (I) contains both a basic moiety, such as an amine or a pyridine or imidazole ring, and an acidic moiety, such as a carboxylic acid. Pharmaceutically acceptable (*i.e.*, non-toxic, physiologically acceptable) salts are preferred, such as, for example, acceptable metal and amine salts in which the cation does not contribute significantly to the toxicity or biological activity of the salt. However, other salts may be useful,  
15   *e.g.*, in isolation or purification steps which may be employed during preparation, and thus, are contemplated within the scope of the invention. Salts of the compounds of the formula (I) may be formed, for example, by reacting a compound of the formula (I) with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

20           Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates,  
25   hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates,  
30   phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts; alkaline earth metal salts such as calcium and magnesium salts; barium, zinc, and aluminum salts; salts with organic bases (for example, organic amines) such as trialkylamines such as triethylamine, procaine, 5 dibenzylamine, N-benzyl- $\beta$ -phenethylamine, 1-phenamine, N,N'-dibenzylethylene-diamine, dehydroabietylamine, N-ethylpiperidine, benzylamine, dicyclohexylamine or similar pharmaceutically acceptable amines and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quaternized with agents such as lower alkyl halides (*e.g.*, methyl, ethyl, propyl, and butyl 10 chlorides, bromides and iodides), dialkyl sulfates (*e.g.*, dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (*e.g.*, decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (*e.g.*, benzyl and phenethyl bromides), and others. Preferred salts include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate or nitrate salts.

15 Compounds of the formula (I) and salts thereof may exist in their tautomeric form, in which hydrogen atoms are transposed to other parts of the molecules and the chemical bonds between the atoms of the molecules are consequently rearranged. It should be understood that the all tautomeric forms, insofar as they may exist, are included within the invention. Additionally, inventive compounds may have *trans* and 20 *cis* isomers and may contain one or more chiral centers, therefore existing in enantiomeric and diastereomeric forms. The invention includes all such isomers, as well as mixtures of *cis* and *trans* isomers, mixtures of diastereomers and racemic mixtures of enantiomers (optical isomers). When no specific mention is made of the configuration (*cis*, *trans* or R or S) of a compound (or of an asymmetric carbon), then 25 any one of the isomers or a mixture of more than one isomer is intended. The processes for preparation can use racemates, enantiomers or diastereomers as starting materials. When enantiomeric or diastereomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional crystallization. The inventive compounds may be in the free or hydrate form.

30 Compounds of the Formula (I) may also have prodrug forms. Any compound that will be converted in vivo to provide the bioactive agent (*i.e.*, the compound for formula I) is a prodrug within the scope and spirit of the invention. Prodrugs and

solvates of the inventive compounds are also contemplated. For example, compounds containing a carboxy group can form physiologically hydrolyzable esters which serve as prodrugs by being hydrolyzed in the body to yield formula (I) compounds *per se*. Such prodrugs are preferably administered orally since hydrolysis in many instances occurs principally under the influence of the digestive enzymes. Parenteral administration may be used where the ester *per se* is active, or in those instances where hydrolysis occurs in the blood. Examples of physiologically hydrolyzable esters of compounds of formula (I) include C<sub>1-6</sub>alkylbenzyl, 4-methoxybenzyl, indanyl, phthalyl, methoxymethyl, C<sub>1-6</sub>alkanoyloxy-C<sub>1-6</sub>alkyl, *e.g.* acetoxymethyl, pivaloyloxymethyl or propionyloxymethyl, C<sub>1-6</sub>alkoxycarbonyloxy-C<sub>1-6</sub>alkyl, *e.g.* methoxycarbonyl-oxymethyl or ethoxycarbonyloxymethyl, glycyloxymethyl, phenylglycyloxymethyl, (5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl and other well known physiologically hydrolyzable esters used, for example, in the penicillin and cephalosporin arts. Such esters may be prepared by conventional techniques known in the art.


For further examples of prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, pp. 309-396, edited by K. Widder, et al. (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, pp. 113-191 (1991); and
- c) H. Bundgaard, Advanced Drug Delivery Reviews, Vol. 8, pp. 1-38 (1992), each of which is incorporated herein by reference.

It should further be understood that solvates (*e.g.*, hydrates) of the compounds of Formula (I) are also within the scope of the present invention. Methods of solvation are generally known in the art.

**Preferred Compounds**

Preferred compounds are those having formula (I),



- 5 their enantiomers, diastereomers, and pharmaceutically-acceptable salts, hydrates, and prodrug thereof, in which:

K is O;

Q is C<sub>1-4</sub>alkylene;

Ar is optionally-substituted phenyl;

- 10 J<sub>1</sub> is a bond, -N(R<sub>5</sub>)-, or -C(R<sub>6a</sub>R<sub>7a</sub>)-;

J<sub>2</sub> is -N(R<sub>5</sub>)- or -C(R<sub>6b</sub>R<sub>7b</sub>)-;

J<sub>3</sub> is -N(R<sub>5</sub>)- or -C(R<sub>6c</sub>R<sub>7c</sub>)-;

provided, however, that only one of J<sub>1</sub>, J<sub>2</sub> and J<sub>3</sub> may be -N(R<sub>5</sub>)-, so that ring A is a five-to-six membered cycloalkyl or heterocyclo ring having from 0 to 2

- 15 heteroatoms;

Y is N or C(R<sub>8</sub>);

Z is N or C(R<sub>9</sub>);

R<sub>1</sub> is selected from hydrogen, C<sub>1-6</sub>alkyl, -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), or C<sub>1-6</sub>alkyl substituted with one to two of hydroxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

R<sub>2</sub> and R<sub>3</sub> are selected from halogen, (C<sub>1-4</sub>)alkyl, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, nitro, phenyloxy, benzyloxy, and phenylthio;

$R_5$  is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cyano,  $-OR_{14}$ ,  $-NR_{14}R_{15}$ ,  $-C(=O)R_{14}$ ,  $-CO_2R_{14}$ ,  $-C(=O)NR_{14}R_{15}$ ,  $-S(O)_pR_{15a}$ ,  $-SO_2NR_{14}R_{15}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl;

5  $R_{6a}$ ,  $R_{6b}$ ,  $R_{6c}$ ,  $R_{7a}$ ,  $R_{7b}$ ,  $R_{7c}$  and  $R_8$  are independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, nitro, cyano,  $-SR_{16}$ ,  $-OR_{16}$ ,  $-NR_{16}R_{17}$ ,  $-C(=O)R_{16}$ ,  $-CO_2R_{16}$ ,  $-C(=O)NR_{16}R_{17}$ ,  $-NR_{16}C(=O)R_{17}$ ,  $-NR_{16}C(=O)OR_{17}$ ,  $-S(O)_qR_{17a}$ ,  $-NR_{16}SO_2R_{17a}$ ,  $-SO_2NR_{16}R_{17}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl; or  $R_{6a}$  with  $R_{7a}$ , or  $R_{6b}$  with  $R_{7b}$ , or  $R_{6c}$  with  $R_{7c}$  are taken together to form a keto group  $(=O)$ ;

10  $R_9$  is selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl,  $-SR_{18}$ ,  $-OR_{18}$ ,  $-NR_{18}R_{19}$ ,  $-CO_2R_{18}$ ,  $-C(=O)R_{18}$ ,  $-C(=O)NR_{18}R_{19}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl;


$R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$ ,  $R_{18}$ , and  $R_{19}$  (i) are selected independently of each other from hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo; or (ii) any two of  $R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$ ,  $R_{18}$ , and  $R_{19}$  when attached to the same nitrogen atom (as in  $NR_{14}R_{15}$ ,  $NR_{16}R_{17}$ , or  $NR_{18}R_{19}$ ) may be taken together to form a heteroaryl or heterocyclo ring, with the remainder of  $R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$ ,  $R_{18}$ , and  $R_{19}$  being selected independently from hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo;

20  $R_{15a}$ , and  $R_{17a}$  are independently selected from alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo;

$p$  is 1, 2, or 3; and

$q$  is 1, 2, or 3.

25 In compounds of formula (I), preferably the groups Q-Ar together form:



wherein  $R_{20a}$  and  $R_{20b}$  are independently selected from halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4})$ alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4})$ alkyl,  $-C(=O)(C_{1-4})$ alkyl,  $-C(=O)NH(CH_2)_rCO_2H$ ,  $-C(=O)NH(CH_2)_rCO_2(C_{1-4})$ alkyl, and  $S(O)_2(C_{1-4})$ alkyl; or from phenyl, benzyl, phenyloxy, benzyloxy and  
 5 heteroaryl in turn optionally substituted with one to two of halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4})$ alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-NH_2$ ,  $-NH(C_{1-4})$ alkyl,  $-N(C_{1-4})$ alkyl<sub>2</sub>,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4})$ alkyl, and/or  $-C(=O)(C_{1-4})$ alkyl; or alternatively, two  $R_{20b}$  groups join together with each other or one  $R_{20b}$  joins together with  $R_{20a}$  to form a fused benzo ring;  $n$  is 0, 1, or 2; and  $r$  is 1, 2, 3, or 4.

10

More preferred are compounds where Q-Ar form para-substituted benzyl, more preferably para-cyano-benzyl or para-bromo-benzyl.

In compounds of formula (I), preferably  $J_1$  is a bond or  $-CHR_{6a}-$ ;  
 15  $J_2$  is  $-CHR_{6b}-$ ;  $J_3$  is  $-CHR_{6c}-$ ; and Y is  $C(R_8)$ , wherein  $R_{6a}$ ,  $R_{6b}$ ,  $R_{6c}$  and  $R_8$  are independently selected from hydrogen, halogen, cyano,  $-S(alkyl)$ ,  $-S(phenyl)$ ,  $-O(alkyl)$ , phenyloxy, benzyloxy,  $-O(C_{1-6})$ alkyl,  $-C(=O)H$ ,  $-C(=O)(C_{1-6})$ alkyl,  $-CO_2H$ ,  $-CO_2(C_{1-6})$ alkyl,  $-C(=O)NH_2$ ,  $-C(=O)NH_2$ ,  $-C(=O)NH(C_{1-4})$ alkyl,  $-C(=O)N(C_{1-4})$ alkyl<sub>2</sub>,  $-NH_2$ ,  $-NH(C_{1-4})$ alkyl,  $-N(C_{1-4})$ alkyl<sub>2</sub>, or from  $C_{1-4}$ alkyl, phenyl, four to seven  
 20 membered heterocyclo,  $C_{3-7}$ cycloalkyl, and five to six membered heteroaryl, each of which in turn is optionally substituted with one to two groups selected from halogen, cyano,  $C_{1-4}$ alkyl, hydroxy, trifluoromethyl, trifluoromethoxy,  $-O(C_{1-6})$ alkyl,  $-C(=O)H$ ,  $-C(=O)(C_{1-6})$ alkyl,  $-CO_2H$ ,  $-CO_2(C_{1-6})$ alkyl,  $-C(=O)NH_2$ ,  $-C(=O)NH_2$ ,  $-C(=O)NH(C_{1-4})$ alkyl,  $-C(=O)N(C_{1-4})$ alkyl<sub>2</sub>,  $-NH_2$ ,  $-NH(C_{1-4})$ alkyl,  $-N(C_{1-4})$ alkyl<sub>2</sub>, hydroxy( $C_{1-4})$ alkyl, methoxy( $C_{1-4})$ alkyl, ethoxy( $C_{1-4})$ alkyl, amino( $C_{1-4})$ alkyl, and halo( $C_{1-4})$ alkyl.  
 25


More preferred are compounds where  $J_1$ ,  $J_2$ , and  $J_3$  are each  $-CH_2-$  and Y is CH. Preferably, Z also is CH.

30 In compounds of formula (I), preferably  $R_1$  is hydrogen,  $C_{1-6}$ alkyl,  $-C(=O)(C_{1-6})$ alkyl, or  $C_{1-6}$ alkyl substituted with one of  $-C(=O)H$ ,  $-C(=O)(C_{1-6})$ alkyl,  $-CO_2H$ , or -

CO<sub>2</sub>(C<sub>1-6</sub>alkyl). More preferred are compounds where R<sub>1</sub> is hydrogen, methyl, and ethyl, and most preferred are compounds where R<sub>1</sub> is methyl.

In compounds of formula (I), preferably R<sub>2</sub> and R<sub>3</sub> are both halogen, and more preferably, R<sub>2</sub> and R<sub>3</sub> are both chloro.

Further preferred compounds are those having the formula (Ia),



and enantiomers, diastereomers, and pharmaceutically-acceptable salts, hydrates, solvates, and prodrug thereof, in which:

Q is  $-(CH_2)_t-$

Ar is phenyl optionally substituted with one to two R<sub>20</sub>;

Y is N or C(R<sub>3</sub>);

Z is N or C(R<sub>9</sub>);

R<sub>1</sub> is selected from hydrogen, C<sub>1-6</sub>alkyl, -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), or C<sub>1-6</sub>alkyl substituted with one to two of hydroxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

R<sub>2</sub> and R<sub>3</sub> are independently selected from halogen, (C<sub>1-4</sub>)alkyl, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, nitro, phenyloxy, benzyloxy, and phenylthio;

R<sub>6</sub> and R<sub>8</sub> at each occurrence are independently selected from hydrogen, halogen, cyano, -S(alkyl), -S(phenyl), -O(alkyl), phenyloxy, benzyloxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -

C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)<sub>2</sub>, or from C<sub>1-4</sub>alkyl, phenyl, four to seven membered heterocyclo, C<sub>3-7</sub>cycloalkyl, and five to six membered heteroaryl, each of which in turn is optionally substituted with one to two groups selected from  
 5 halogen, cyano, C<sub>1-4</sub>alkyl, hydroxy, trifluoromethyl, trifluoromethoxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)<sub>2</sub>, hydroxy(C<sub>1-4</sub>)alkyl, methoxy(C<sub>1-4</sub>)alkyl, ethoxy(C<sub>1-4</sub>)alkyl, amino(C<sub>1-4</sub>)alkyl, and halo(C<sub>1-4</sub>)alkyl; and/or two R<sub>6</sub> groups taken  
 10 together form keto (=O);

R<sub>9</sub> is selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl, -S(alkyl), -O(alkyl), S(phenyl), -O(alkyl), phenyloxy, benzyloxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>,  
 15 -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

R<sub>20</sub> at each occurrence is selected from halogen, C<sub>1-6</sub>alkyl, hydroxy, (C<sub>1-4</sub>)alkoxy, (C<sub>1-4</sub>)alkylthio, cyano, nitro, -CO<sub>2</sub>H, -C(=O)H, -CO<sub>2</sub>(C<sub>1-4</sub>)alkyl, -C(=O)(C<sub>1-4</sub>)alkyl, -C(=O)NH(CH<sub>2</sub>)<sub>r</sub>CO<sub>2</sub>H, -C(=O)NH(CH<sub>2</sub>)<sub>r</sub>CO<sub>2</sub>(C<sub>1-4</sub>alkyl), S(O)<sub>2</sub>(C<sub>1-4</sub>alkyl), phenyl, benzyl, phenyloxy, benzyloxy, five to six membered  
 20 heteroaryl, C<sub>3-7</sub>cycloalkyl, and four to seven membered heterocyclo, wherein each of the alkyl, alkoxy, and cyclic groups in turn are optionally substituted with one to three of R<sub>24</sub>;

R<sub>24</sub> is selected from halogen, cyano, nitro, C<sub>1-4</sub>alkyl, hydroxy, trifluoromethyl, trifluoromethoxy, -O(C<sub>1-4</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)<sub>2</sub>, hydroxy(C<sub>1-4</sub>)alkyl, methoxy(C<sub>1-4</sub>)alkyl, ethoxy(C<sub>1-4</sub>)alkyl, amino(C<sub>1-4</sub>)alkyl, and halo(C<sub>1-4</sub>)alkyl;  
 25


*m* is 0 or 1;

*r* and *s* are 0, 1, 2, 3 or 4; and

30 *t* is 1 or 2.



More preferred are compounds having the formula (Ib),




- 5 in which  $R_1$ ,  $R_2$ ,  $R_3$ ,  $Y$  and  $Z$  are selected as recited above for compounds of formula (I), and  $R_{20a}$  and  $R_{20b}$  are independently selected from halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4})$ alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4})$ alkyl,  $-C(=O)(C_{1-4})$ alkyl,  $-C(=O)NH(CH_2)_rCO_2H$ ,  $-C(=O)NH(CH_2)_rCO_2(C_{1-4})$ alkyl, and  $S(O)_2(C_{1-4})$ alkyl; or from phenyl, benzyl, phenyloxy, benzyloxy and heteroaryl in turn optionally substituted with one to two of
- 10 halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4})$ alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-NH_2$ ,  $-NH(C_{1-4})$ alkyl,  $-N(C_{1-4})_2$ ,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4})$ alkyl, and/or  $-C(=O)(C_{1-4})$ alkyl; or alternatively, two  $R_{20b}$  groups join together with each other or one  $R_{20b}$  joins together with  $R_{20a}$  to form a fused benzo ring; and  $n$  is 0, 1 or 2.

15

Further preferred compounds are those of formula (Ib), as immediately defined above, wherein  $R_1$  is  $C_{1-4}$  alkyl,  $R_2$  and  $R_3$  are both halogen (more preferably chlorine),  $R_{20a}$  is cyano or halogen, and  $n$  is 0.

20

Most preferred are compounds having the formula,



wherein preferably Z is CH, R<sub>1</sub> is methyl or ethyl, and R<sub>20a</sub> is cyano or halogen.

5


### Methods of Preparation

The compounds of the invention may be prepared by the exemplary processes described in the following reaction Schemes A-L. Exemplary reagents and procedures for these reactions appear hereinafter. Starting materials are commercially available or can be readily prepared by one of ordinary skill in the art. For all of the

10 schemes, the groups Z, K, Q, Ar, Y, J<sub>1</sub>, J<sub>2</sub>, J<sub>3</sub>, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as described herein for a compound of formula (I), unless otherwise indicated. Groups designated generally as R, R', X, and P as well as solvents, temperatures, pressures, starting materials having the desired groups, and other reaction conditions, may be readily selected as appropriate by one of ordinary skill in the art.


15

### Scheme A:




A suitably functionalized amino ketone 1 is reacted with an isocyanate 2a or an isothiocyanate 2b in DCM to yield the ureidoketone 3a or the thioureidoketone 3b, respectively. This intermediate product is then cyclized in an organic solvent such as xylene in the presence of a catalytic amount of base (such as  $K_2CO_3$ ) and molecular sieves at a temperature of 100-140°C to give the hexahydro-benzimidazolone having the formula (Ic) or hexahydro-benzimidazol-thione having the formula (Id).

Compounds of formulae 1, 2a, and 2b are commercially available or can be readily prepared by one skilled in the field as shown in *e.g.*, Demaesle *et al.*, Tetrahedron Lett., Vol. 33 (1992), at pp. 4447-4450; Parcell *et al.*, J. Org. Chem., Vol. 46 (1981), at pp. 5229-5231; or Benati *et al.*, Tetrahedron, Vol. 58 (2002), at pp. 3485-3492.

**Scheme B:**


- A suitably functionalized amino ketone **1** is reacted with an isocyanate **2a** or an isothiocyanate **2b** in EtOAc in the presence of a base (such as Et<sub>3</sub>N, KOH, K<sub>2</sub>CO<sub>3</sub>, KHCO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub> or NaHCO<sub>3</sub>), to give after treating with MeOH the hydroxy bicyclic-imidazolone **4a** or the hydroxy bicyclic-imidazolthione **4b**, respectively. This intermediate product is then dehydrated in an organic solvent (such as toluene) in the presence of a catalytic amount of dehydrating agent (such as PTSA or molecular sieves) at a temperature of 100-140°C to yield the hexahydro-benzimidazolone having the formula (Ic) or hexahydro-benzimidazol-thione having the formula (Id). See, e.g., Bobowski *et al.*, *J. Heterocycl Chem.*, Vol.18 (1981), at p. 1179-1187. Compounds of formulae **1**, **2a**, and **2b** are commercially available or can be readily prepared by one skilled in the field as shown in e.g., Demaesle *et al.*, *Tetrahedron Lett.*, Vol. 33 (1992), at pp. 4447-4450; Parcell *et al.*, *J. Org. Chem.*, Vol. 46 (1981), at pp. 5229-5231; or Benati *et al.*, *Tetrahedron*, Vol. 58 (2002), at pp. 3485-3492.

**Scheme C:**




A suitably functionalized Boc-protected amino ketone **1a** is heated with an aniline **5** in toluene, to yield the imine **6**. This intermediate product is then reacted, after deprotection, with a coupling agent such as triphosgene in an organic solvent (such as DCM) in the presence of base (such as pyridine) to yield the hexahydrobenzimidazolone having the formula (Ie). See, e.g., Rao *et al.*, *Tetrahedron Lett.*, Vol.34 (1993), at p. 4993. Compounds of formulae **1a** and **5** are commercially available or can be readily prepared by one skilled in the field as shown in e.g., Demaesle *et al.*, *Tetrahedron Lett.*, Vol. 33 (1992), at pp. 4447-4450 or Parcell *et al.*, *J. Org. Chem.*, Vol. 46 (1981), at pp. 5229-5231, or Benati *et al.*, *Tetrahedron*, Vol. 58 (2002), at pp. 3485-3492

#### Scheme D:




Treatment of bicyclic imidazolone (Ic) with a reagent such as  $P_2S_5$  or 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's reagent) in an organic solvent such as toluene or dioxane, yields the corresponding hexahydro-benzimidazol-thione having the formula (Id). See, e.g., Rao *et al.*, *Tetrahedron Lett.*, Vol. 34 (1993), at p. 4993.

**Scheme E:**




*N*-acylation of bicyclic imidazolone having the formula (If) can be obtained by treatment with an acylating agent (such as an acyl chloride, an anhydride, a chloroformate or an isocyanate) in an organic solvent (such as THF or acetonitrile) in the presence of a base (such as TEA, DIPEA, DMAP or NaH) to give bicyclic imidazolone having the formula (Ig). *See, e.g.,* Melnyk *et al.*, Tetrahedron, Vol. 48 (1992), at pp. 841-850, and Ross *et al.*, Syn. Commun., Vol. 28 (1998), at p. 3877-3884.

**Scheme F:**




Sulfonylation of hexahydro-benzimidazolone having the formula (If) is obtained by treatment with a sulfonyl chloride in the presence of a base (such as TEA, DIPEA, pyridine or DMAP) in an organic solvent such as toluene to give the desired compounds having the formula (Ih). *See, e.g., Yokoyama et al., Tetrahedron Lett.*, Vol. 39 (1998), at pp. 4847-4850.

**Scheme G:**



Bicyclic imidazolone having the formula (If) can be *N*-alkylated in an aprotic solvent (such as DMF, THF or DMSO) by treatment with one equivalent of a base (such as NaH, NaHMDS, LDA, LiHMDS, KH, KHMDS or *t*BuOK) followed by addition of a suitable alkylating agent (such as an alkyl iodide, alkyl bromide, alkyl chloride, a tosylate or a mesylate) to yield hexahydro-benzimidazolone having the formula (Ii). *See, e.g., Melloni et al., Farmaco*, Vol. 9 (1991), at pp. 1011-1021.


**Scheme H:**




Bromo bicyclic imidazolone having the formula (Ij) can be transformed into biaryl compounds having the formula (Ik) by reaction with an aromatic or

heteroaromatic boronic acid **10** in the presence of a palladium catalyst (such as  $\text{Pd}(\text{PPh}_3)_4$ ) and a base (such as  $\text{K}_2\text{CO}_3$  or  $\text{Na}_2\text{CO}_3$ ) in an appropriate solvent (such as toluene, DMF, DME or water) under conventional Suzuki coupling conditions. See, e.g., Suzuki *et al.*, *Synth. Commun.*, Vol. 11 (1981), at p. 513.

5

**Scheme I:**


Bromo bicyclic imidazolone having the formula (**Ij**) can be transformed into biaryl compounds having the formula (**Ik**) by reaction with an aromatic or heteroaromatic alkyl stannate **11** in the presence of a palladium catalyst (such as  $\text{Pd}(\text{PPh}_3)_4$ ) in an appropriate solvent (such as toluene, DMF, DME or water) under conventional Stille coupling conditions. See, e.g., Yang *et al.*, *Synth. Commun.*, Vol. 22; 12; (1992), at pp. 1757-1762.

15 **Scheme J:**




Bicyclic imidazolone (II) can be debenzylated using for example 1-chloroethyl chloroformate in a solvent such as dichloromethane or dichloroethane to yield the NH derivative having the formula (Im). This compound can be alkylated either by reaction with an alkyl halide  $R_5X$  (for example an alkyl iodide) in a solvent such as acetonitrile or acetone at temperature ranging from room temperature to reflux, or by reaction with an aldehyde in the presence of a reducing agent such as sodium triacetoxy borohydride or sodium cyanoborohydride in a solvent such as acetonitrile or dichloroethane to yield the compound of formula (It).

10 **Scheme K:**



Compounds of formula (Im) can also be acylated with an acyl halide (for example an acyl chloride or acyl bromide) in the presence of an organic base (such as triethylamine or diisopropylethylamine) or an inorganic base (such as sodium carbonate) in a solvent such as dichloromethane at temperature ranging from  $-15^{\circ}\text{C}$  to room temperature to yield the acylated derivative of formula (In). The same compound can also be obtained by reaction with an acid  $\text{RCO}_2\text{H}$  in the presence of a coupling agent such as dicyclohexylcarbodiimide in a solvent such as dichloromethane.

**Scheme L:**

Sulfonamide of formula (Io) can be obtained from compound of formula (Im)

5 by reaction with a sulfonyl halide in the presence of a base (such as triethylamine or sodium carbonate) in a solvent such as dichloromethane or tetrahydrofuran at temperature ranging from  $-15^{\circ}\text{C}$  to room temperature.

**Utility**

10 The compounds and compositions of this invention are antagonists and/or inhibitors of LFA-1, Mac-1, and/or ICAMs. They are useful in treating various inflammatory diseases and disorders associated with the action of LFA-1, Mac-1, and/or ICAMs, particularly LFA-1:ICAM-1. The term "Leukointegrin/ICAM-associated condition" is used herein for ease of reference to refer to those diseases or disorders that are associated with the action or levels of LFA-1, Mac-1 and/or ICAM-1, ICAM-2, or ICAM-3. As used herein, the term "treating" includes prophylactic and therapeutic uses and thus includes the alleviation of symptoms of a

15 Leukointegrin/ICAM-associated condition in a patient, the improvement of an ascertainable measurement associated with such a condition, or the prevention of such a condition or its symptoms. The term "patient" refers to a mammal, preferably a human.

In view of their inhibition activity, the compounds may be used to treat conditions involving the activation, co-stimulation, or infiltration of T-cells and/or leukocytes, including without limitation conditions involving the influx of leukocytes

25 in the skin, peritoneum, synovium, lung, kidney, and heart. The inventive compounds

may be used to treat conditions resulting from a response of the specific or non-specific immune system in a patient.

Leukointegrin/ICAM-associated conditions that may be treated with the inventive compounds include acute or chronic graft vs host reactions (*e.g.*, pancreatic islet allograft); and acute or chronic transplant rejection (*e.g.*, kidney, liver, heart, lung, pancreas, bone marrow, cornea, small bowel, skin allografts, skin homografts, heterografts, and/or cells derived from such organs). Additionally, the compounds may be used to treat inflammatory conditions including, but not limited to, multiple sclerosis, rheumatoid arthritis, psoriatic arthritis, osteoarthritis, osteoporosis, diabetes (*e.g.*, insulin dependent diabetes mellitus or juvenile onset diabetes), cystic fibrosis, inflammatory bowel disease, irritable bowel syndrome, Crohn's disease, ulcerative colitis, Alzheimer's disease, shock, ankylosing spondylitis, gastritis, conjunctivitis, pancreatitis (acute or chronic), multiple organ injury syndrome (*e.g.*, secondary to septicemia or trauma), myocardial infarction, atherosclerosis, stroke, reperfusion injury (*e.g.*, due to cardiopulmonary bypass or kidney dialysis), acute glomerulonephritis, vasculitis, thermal injury (*i.e.*, sunburn), necrotizing enterocolitis, granulocyte transfusion associated syndrome, and/or Sjogren's syndrome.

The inventive compounds may be used to treat inflammatory conditions of the skin including eczema, atopic dermatitis, contact dermatitis, urticaria, scleroderma, psoriasis, and dermatosis with acute inflammatory components.

The compounds also may also be used to treat allergies and respiratory conditions, including asthma, pulmonary fibrosis, allergic rhinitis, oxygen toxicity, emphysema, chronic bronchitis, acute respiratory distress syndrome, and any chronic obstructive pulmonary disease (COPD). The compounds may be used to treat chronic hepatitis infection, including hepatitis B and hepatitis C.

Additionally, the compounds of the invention may be used to treat autoimmune diseases and/or inflammation associated with autoimmune diseases such as organ-tissue autoimmune diseases (*e.g.*, Raynaud's syndrome), autoimmune thyroiditis, uveitis, systemic lupus erythematosus, Addison's disease, autoimmune polyglandular disease (also known as autoimmune polyglandular syndrome), and Grave's disease.

The compounds of this invention also may be used to treat metastases or as an adjunct to minimize toxicity with cytokine therapy in the treatment of cancers.

The compounds of this invention further have utility in treating hypogonadism, frailty, sexual dysfunction, wasting, such as wasting syndromes associated with cancer and AIDS, and anemia. The compounds further have utility in treating cancers, including but not limited to cancers of the breast, brain, skin, ovary, endometrium, bladder, prostate, lung, colon, lymphatic system, liver and kidney. The inventive compounds are useful for conditions such as hirsutism, acne, seborrhea, alopecia, fibroids, hyperpilosity, cachexia, polycystic ovarian syndrome, anorexia, contraception, drug withdrawal syndrome, pregnancy termination, and benign prostate hypertrophy. The compounds are further useful as antiangiogenic agents. Additionally, the compounds may be useful as inhibitors of protein prenyltransferases, particularly farnesyltransferase and the prenylation of the oncogene protein Ras. As such, the inventive compounds may potentially be useful for treating and/or preventing the diseases and disorders referred to in WO 01/45704, incorporated herein by reference.

When used as anti-inflammatory agents, the compounds may be administered prior to the onset of, at, or after the initiation of inflammation. When used prophylactically, the compounds are preferably provided in advance of any inflammatory response or symptom (for example, prior to, at, or shortly after the time of an organ or tissue transplant but in advance of any symptoms of organ rejection). Administration of the compounds may prevent or attenuate inflammatory responses or symptoms.

The present invention thus provides methods for treating such conditions as those listed above, comprising administering to a patient in need thereof an effective amount of at least one compound of formula (I) or a salt thereof. Other therapeutic agents such as those described below may be employed in combination with the compounds of formula (I). In the methods of the present invention, such other therapeutic agent(s) may be administered prior to, simultaneously with, or following the administration of the inventive compounds.

The present invention also provides pharmaceutical compositions capable of treating the above-referenced diseases and disorders. The inventive compositions may contain other therapeutic agents and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.), according to techniques such as those well known in the art of pharmaceutical formulation.

The compounds of formula (I) may be administered by any means suitable for the condition to be treated, which may depend on the need for site-specific treatment or quantity of drug to be delivered. Topical administration is generally preferred for skin-related diseases, and systematic treatment preferred for cancerous or pre-cancerous conditions, although other modes of delivery are contemplated. For example, the compounds may be delivered orally, such as in the form of tablets, capsules, granules, powders, or liquid formulations including syrups; topically, such as in the form of solutions, suspensions, gels or ointments; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (*e.g.*, as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; rectally such as in the form of suppositories; or liposomally. Dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents may be administered. The compounds may be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved with suitable pharmaceutical compositions or particularly in the case of extended release, with devices such as subcutaneous implants or osmotic pumps.

Exemplary compositions for topical administration include a topical carrier such as PLASTIBASE® (mineral oil gelled with polyethylene).

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium

phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The inventive compounds may also be orally delivered by sublingual and/or buccal administration, *e.g.*, with molded, compressed, or freeze-dried tablets. Exemplary  
5 compositions may include fast-dissolving diluents such as mannitol, lactose, sucrose, and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (AVICEL®) or polyethylene glycols (PEG); an excipient to aid mucosal adhesion such as hydroxypropyl cellulose (HPC), hydroxypropyl methyl cellulose (HPMC), sodium carboxymethyl cellulose (SCMC),  
10 and/or maleic anhydride copolymer (*e.g.*, GANTREZ®); and agents to control release such as polyacrylic copolymer (*e.g.*, CARBOPOL 934®). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal administration via aerosol or inhalation  
15 include solutions which may contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance absorption and/or bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or suspensions which may contain, for example, suitable non-toxic,  
20 parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Exemplary compositions for rectal administration include suppositories  
25 which may contain, for example, suitable non-irritating excipients, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures but liquefy and/or dissolve in the rectal cavity to release the drug.

The effective amount of a compound of the present invention may be determined by one of ordinary skill in the art, and includes exemplary dosage amounts  
30 for a patient of from about 0.05 to 100 mg/kg of body weight of active compound per day, which may be administered in a single dose or in the form of individual divided

doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors, including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, 5 age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and the particular condition sought to be treated and its severity. Preferred subjects for treatment include animals, most preferably mammalian species such as humans, and domestic animals such as dogs, cats, horses, and the like, subject to Leukointegrin/ICAM associated conditions and/or subject to any of the above-referenced diseases and disorders. 10

The inventive compounds and compositions may be employed alone or in combination with each other and/or other suitable therapeutic agents useful in treating diseases and disorders referenced above, for example, where the second drug has the same or different mechanism of action than the present compounds. Exemplary of 15 such other therapeutic agents include anti-inflammatory agents, antibiotics, anti-viral agents, anti-oxidants, and agents used to treat respiratory conditions such as COPD and asthma.

Examples of suitable other anti-inflammatory agents with which the inventive compounds may be used include aspirin, cromolyn, nedocromil, theophylline, 20 zileuton, zafirlukast, montelukast, pranlukast, indomethacin, and lipoxygenase inhibitors; non-steroidal antiinflammatory drugs (NSAIDs) (such as ibuprofen and naproxin); TNF- $\alpha$  inhibitors (such as tenidap and rapamycin or derivatives thereof), or TNF- $\alpha$  antagonists (*e.g.*, infliximab, Enbrel®, D2E7, OR1384), cytokine modulators (*e.g.* TNF-alpha converting enzyme [TACE] inhibitors, Interleukin-1 25 converting enzyme (ICE) inhibitors, Interleukin-1 receptor antagonists), prednisone, dexamethasone, cyclooxygenase inhibitors (*i.e.*, COX-1 and/or COX-2 inhibitors such as Naproxen®, Celebrex®, or Vioxx®), CTLA4-Ig agonists/antagonists (LEA29Y), CD40 ligand antagonists, IMPDH inhibitors (such as mycophenolate [CellCept®] and merimepodib), methotrexate (FK506), integrin antagonists (*e.g.*, alpha-4 beta-1, 30 alpha-V-beta-3), cell adhesion inhibitors, interferon gamma antagonists, prostaglandin synthesis inhibitors, budesonide, clofazimine, CNI-1493, CD4 antagonists (*e.g.*, priliximab), p38 mitogen-activated protein kinase inhibitors, protein tyrosine kinase

(PTK) inhibitors, IKK inhibitors, therapies for the treatment of irritable bowel syndrome (*e.g.*, Zelmac®, Zelnorm®, and Maxi-K® openers such as those disclosed in U.S. Patent No. 6,184,231 B1), or NF- $\kappa$ B inhibitors (such calphostin, CSAIDs, and quinoxalines as disclosed in US Pat. No. 4,200,750); disassociated steroids;

5 chemokine receptor modulators (including CCR1, CCR2, CCR3, CCR4, and CXCR2 receptor antagonists); secretory and cytosolic phospholipase A2 inhibitors, glucocorticoids, salicylates, nitric oxide, and other immunosuppressants; and nuclear translocation inhibitors, such as deoxyspergualin (DSG).

The inventive compounds may be used in combination with other agents used

10 to treat respiratory conditions such as asthma, COPD, and allergic rhinitis, such as  $\beta$ -adrenergic agonists (such as albuterol, terbutaline, formoterol, salbutamol, salmeterol, bitolterol, pilbuterol, and fenoterol); corticosteroids (such as beclomethasone, triamcinolone, budesonide, fluticasone, flunisolide, dexamethasone, prednisone, and dexamethasone); leukotriene antagonists (*e.g.*, Accolate [Zafirlukast®], and Singulair

15 [Montelukast®]); Muscarinic M3 cholinergic antagonists (*e.g.*, Spiriva®), PDE 4 inhibitors (*e.g.* rolipram, cilomilast [Ariflo®], piclamilast, or roflumilast), histamine H<sub>1</sub> antagonists, Allegra® (fexofenadine), Claritin® (loratidine), and/or Clarinex® (desloratidine).

Examples of suitable antiviral agents for use with the inventive compounds

20 include nucleoside-based inhibitors, protease-based inhibitors, and viral-assembly inhibitors.

Examples of suitable anti-osteoporosis agents for use in combination with the compounds of the present invention include alendronate, risedronate, PTH, PTH fragment, raloxifene, calcitonin, RANK ligand antagonists, calcium sensing receptor

25 antagonists, TRAP inhibitors, selective estrogen receptor modulators (SERM) and AP-1 inhibitors.

Examples of suitable anti-oxidants for use in combination with the compounds of the present invention include lipid peroxidation inhibitors such as probucol, BO-653, Vitamin A, Vitamin E, AGI-1067, and  $\alpha$ -lipoic acid.



The inventive compounds also may be used in combination with anti-diabetic agents, such as biguanides (*e.g.* metformin), glucosidase inhibitors (*e.g.* acarbose), insulins (including insulin secretagogues or insulin sensitizers), meglitinides (*e.g.* repaglinide), sulfonylureas (*e.g.*, glimepiride, glyburide and glipizide),  
5 biguanide/glyburide combinations (*e.g.*, glucovance), thiozolidinediones (*e.g.* troglitazone, rosiglitazone and pioglitazone), PPAR-alpha agonists, PPAR-gamma agonists, PPAR alpha/gamma dual agonists, SGLT2 inhibitors, inhibitors of fatty acid binding protein (aP2) such as those disclosed in U.S. Serial No. 09/519,079 filed March 6, 2000 and assigned to the present assignee, glucagon-like peptide-1 (GLP-1),  
10 glucagon phosphorylase, and dipeptidyl peptidase IV (DP4) inhibitors.

In addition, the compounds may be used with agents that increase the levels of cAMP or cGMP in cells for a therapeutic benefit. For example, the compounds of the invention may have advantageous effects when used in combination with phosphodiesterase inhibitors, including PDE1 inhibitors (such as those described in  
15 Journal of Medicinal Chemistry, Vol. 40, pp. 2196-2210 [1997]), PDE2 inhibitors, PDE3 inhibitors (such as revizinone, pimobendan, or olprinone), PDE4 inhibitors (referenced above), PDE7 inhibitors, or other PDE inhibitors such as dipyridamole, cilostazol, sildenafil, denbutyline, theophylline (1,2-dimethylxanthine), ARIFLO™ (*i.e.*, cis-4-cyano-4-[3-(cyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic  
20 acid), arofyline, C-11294A, CDC-801, BAY-19-8004, cipamfylline, SCH351591, YM-976, PD-189659, mesiopram, pumafentrine, CDC-998, IC-485, and KW-4490.

In view of their usefulness in treating ischemia, the inventive compounds may be used in combination with agents for inhibiting  $F_1F_0$ -ATPase, including efrapeptin, oligomycin, autovertin B, azide, and compounds described in US patent application  
25 Serial No. 60/339,108, filed December 10, 2001 and assigned to the present assignee; -alpha- or beta- adrenergic blockers (such as propranolol, nadolol, carvedilol, and prazosin), antianginal agents such as nitrates, for example, sodium nitrates, nitroglycerin, isosorbide mononitrate, isosorbide dinitrate, and nitrovasodilators; antiarrhythmic agents including Class I agents (such as propafenone); Class II agents  
30 (propranolol); Class III agents (such as sotalol, dofetilide, amiodarone, azimilide and ibutilide); Class IV agents (such as diltiazem and verapamil);  $K^+$  channel modulators such as  $I_{ACh}$  inhibitors and inhibitors of the  $K_v1$  subfamily of  $K^+$  channel openers such

as  $I_{K_{ATP}}$  inhibitors (*e.g.*, compounds disclosed in U.S. Application Serial No. 09/729,731, filed December 5, 2000); and gap-junction modulators such as connexions; anticoagulant or antithrombotic agents including aspirin, warfarin, ximelagran, low molecular weight heparins (such as lovenox, enoxaparin, and  
5 dalteparin), anti-platelet agents such as GPIIb/GPIIIa blockers, (*e.g.*, abciximab, eptifibatide, and tirofiban), thromboxane receptor antagonists (*e.g.*, ifetroban),  $P2Y_1$  and  $P2Y_{12}$  antagonists (*e.g.*, clopidogrel, ticlopidine, CS-747, and aspirin/clopidogrel combinations), and Factor Xa inhibitors (*e.g.*, fondaparinux); and diuretics such as sodium-hydrogen exchange inhibitors, chlorothiazide, hydrochlorothiazide,  
10 flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide, benzthiazide, ethacrynic acid, triacryfen, chlorthalidone, furosemide, musolimine, bumetanide, triamterene, and amiloride.

The inventive compounds may also be useful in combination with antiangiogenic agents, such as compounds that are inhibitors of VEGF receptors, or in  
15 conjunction with antitumor agents such as paclitaxel, adriamycin, epithilones, cisplatin, and carboplatin. Examples of anticancer and other cytotoxic agents that may be used in combination with the inventive compounds include the following: epothilone derivatives as found in German Patent No. 4138042.8; WO 97/19086, WO 98/22461, WO 98/25929, WO 98/38192, WO 99/01124, WO 99/02224, WO  
20 99/02514, WO 99/03848, WO 99/07692, WO 99/27890, WO 99/28324, WO 99/43653, WO 99/54330, WO 99/54318, WO 99/54319, WO 99/65913, WO 99/67252, WO 99/67253 and WO 00/00485; cyclin dependent kinase inhibitors as found in WO 99/24416; and prenyl-protein transferase inhibitors as found in WO 97/30992 and WO 98/54966.

25       The combination of the inventive compounds with other therapeutic agents may prove to have additive and synergistic effects. The combination may be advantageous to increase the efficacy of the administration or decrease the dosage to reduce possible side-effects.

      The above other therapeutic agents, when employed in combination with  
30 the compounds of the present invention, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art. In the methods of the present invention, such other

therapeutic agent(s) may be administered prior to, simultaneously with, or following the administration of the inventive compounds.

Compounds of formula (I), including the compounds described in the examples hereof, have been tested in assay(s) described below and have shown a measurable level of activity as inhibitors of LFA-1 and/or ICAM-1.

### Assays

#### H1-HeLa Adhesion assay

H1-Hela cells were released from their growth flask using versene (Gibco, Grand Island, NY). Following centrifugation, the cells were resuspended in growth medium: DMEM (Gibco), 10% fetal calf serum (Hyclone, Logan, UT), 1% Pen-Strep (Gibco), and 1% L-glutamine (Gibco) and plated for growth at 5,000 cells/well in a 96-well plate.

The next day, HSB-2 cells were divided to  $2 \times 10^5$ /ml in growth medium: RPMI 1640 (Gibco), 10% FCS, 1% Pen-Strep, and 1% L-glutamine. The next day (day #3), the cells were centrifuged at 534xG for 8 minutes, washed, and resuspended in HBSS at  $5 \times 10^7$ /ml. Calcein-AM, 10  $\mu$ M (Molecular Probes, Eugene, OR) and 100 nM phorbol myristate acetate (SIGMA, St. Louis, MO) were added to the labeling and activation mix. Following incubation at 37°C for 30 minutes, ten ml of HBSS was added and the cells centrifuged as above. The cell pellet was then resuspended and counted.

While the HSB-2 cells were labeling, the medium was aspirated from the H1-HeLa cells and the plates washed once with HBSS, followed by the addition of 50  $\mu$ l of HBSS. An additional 50  $\mu$ l of HBSS containing compound solution, DMSO, or anti-CD18 antibody was then added to each well. To the H1-HeLa cells were added 200,000 HSB-2 cells/well in 100  $\mu$ l, followed by incubation in the dark for 30 minutes. The wells were then washed three times to remove the unbound cells. A fluorescence plate reader was then used to determine the number of bound HSB-2 cells. The percent inhibition due to the compound was calculated using the vehicle control as 0% inhibition and the antibody blocked adhesion as 100% inhibition.

HUVEC adhesion assay

On day 1, human umbilical vein endothelial cells (HUVEC) (passage 3, Clonetics, San Diego, CA) were placed into a T-75 flask containing EGM bulletkit media (Clonetics) for growth.

5        When the HUVEC were 90% confluent (typically day 4), 96-well tissue culture plates were coated with 100  $\mu$ l/well of 2.5  $\mu$ g/ml mouse Type IV collagen (Trevigen) diluted in 0.1 M acetic acid. Following incubation for at least three hours, the collagen was removed and the plate washed three times with HBSS (Gibco). The HUVEC flask was trypsinized, and HUVEC were plated on the collagen coated wells  
10        at 1250 cells/200  $\mu$ l/well for use four days later. Twenty hours prior to use, the medium was removed and cells were stimulated with 200  $\mu$ l of 10nM phorbol myristate acetate (PMA, Sigma, St. Louis, MO) in EGM. When the cells were 90% confluent (typically day 8), the PMA-containing medium was removed, the wells were washed with HBSS, and 50  $\mu$ l of HBSS was added to the wells. An additional  
15        50  $\mu$ l containing compound solution, DMSO or blocking anti-CD18 was then added to each well.

On day 7, HSB-2 cells were then divided to  $2 \times 10^5$ /ml in RPMI 1640 (Gibco), 10% FCS (Hyclone, Logan, UT), 1% Pen-Strep (Gibco), and 1% L-glutamine (Gibco). The following day, the cells were centrifuged at 534xG for 8 minutes,  
20        washed, and resuspended in HBSS at  $5 \times 10^7$ /ml. For activation and labeling, calcein-AM, 10  $\mu$ M (Molecular Probes, Eugene, OR) and 100 nM phorbol myristate acetate (Sigma, St. Louis, MO) were added and the cells incubated at 37°C for 30 minutes. Following the addition of ten ml of HBSS, the cells were centrifuged, resuspended, and counted.

25        To the HUVEC cells were added 200,000 labeled and activated HSB-2 cells/well in 100  $\mu$ l, followed by incubation in the dark for 30 minutes. To remove unbound cells, the wells were washed three times with HBSS. A fluorescence plate reader was used to determine the number of HSB-2 cells bound. The percent inhibition due to the compound was calculated with the vehicle control set at 0%  
30        inhibition and the antibody-blocked adhesion set at 100% inhibition.

### EXAMPLES

The following Examples illustrate embodiments of the inventive compounds and starting materials, and are not intended to limit the scope of the claims. For ease of reference, the following abbreviations are used herein:

5

#### Abbreviations

- bp = boiling point  
CH<sub>3</sub>CN = acetonitrile  
DCC = dicyclohexylcarbodiimide  
10 DCE = dichloroethane  
DCM = dichloromethane  
DMAP = 4-dimethylaminopyridine  
DIPEA or DIEA = *N,N*-diisopropylethylamine  
DMF = dimethyl formamide  
15 EDCI = 1-3-dimethylaminopropyl-3-ethylcarbodiimide  
HOBT = 1-hydroxybenzotriazole  
EtOAc = ethyl acetate  
EtOH = ethanol  
g = gram(s)  
20 HCl = hydrochloric acid  
KOH = potassium hydroxide  
K<sub>2</sub>CO<sub>3</sub> = potassium carbonate  
l = liter  
LiAlH<sub>4</sub> = lithium aluminum hydride  
25 MeCN = acetonitrile  
MeOH = methanol  
MgSO<sub>4</sub> = magnesium sulfate  
NaH = sodium hydride  
Na<sub>2</sub>SO<sub>4</sub> = sodium sulfate  
30 NaOH = sodium hydroxide  
NMP = 1-methyl-2-pyrrolidinone  
PBr<sub>3</sub> = phosphorus tribromide  
(Ph<sub>3</sub>P)<sub>4</sub>Pd = tetrakis(triphenylphosphine)palladium(0)  
SOCl<sub>2</sub> = thionyl chloride  
35 tBuOMe = t-Butyl methyl ether  
TEA = triethylamine  
mg = milligram(s)  
ml = milliliter  
μl = microliter

mmol = millimole

$\mu$ mol = micromole


mol = mole

mp = melting point

5 RT = room temperature

### Preparation 1

#### 4-[(2-oxocyclohexyl)methyl]-benzonitrile




10

NaI (1g) was added to a solution of 4-Bromomethyl benzonitrile (12.9 g, 0.066 mol) and 1-(pyrrolidino-1-cyclohexene (10 g, 0.066 mol) in EtOH (150 ml). After 1 night at RT, the reaction mixture was poured into water (500 ml) and  
15 extracted twice with EtOAc. The organic phase was washed twice with water, dried over magnesium sulfate and concentrated to yield 4-[(2-oxocyclohexyl)methyl]-benzonitrile (7.06 g) as a yellow oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 7.60 (2H, d), 7.30 (2H, d), 3.25 (1H, dd), 2.7-2.2 (4H, m), 2.2-1.2 (6H, m).

20

### Preparation 2

#### 4-[[2-(dimethylhydrazono)cyclohexyl]methyl]-benzonitrile




25 A solution of 4-[(2-oxocyclohexyl)methyl]-benzonitrile (7 g) (0.033 mol) and anhydrous N,N-dimethylhydrazine (9.2-ml) (3,7 eq.) in EtOH (100 ml) was refluxed for 8 h. After cooling, the reaction mixture was concentrated to dryness. The residue

was partitioned between tBuOMe and water, the organic phase washed twice with water, dried over magnesium sulfate, and concentrated to yield 4-[[2-(dimethylhydrazono)cyclohexyl]methyl]-benzonitrile (6.1 g) as a orange oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.30 (2H, d), 3.15 (1H, dd), 2.9-2.15 (6H, m), 2.3 (6H, s), 1.8-1.3 (4H, m).

### Preparation 3


#### **2-[(4-bromophenyl)methyl]-cyclohexanone, dimethylhydrazone**



Anhydrous N,N-dimethylhydrazine (17 ml) (3,7 eq.) was quickly poured in a solution of 2-[(4-bromophenyl)methyl]-cyclohexanone (16 g) (0.06 mol) in EtOH (200 ml). The reaction mixture was refluxed for 10 h, then concentrated to dryness. The residue was partitioned between tBuOMe and water, the organic phase washed twice with water, dried over magnesium sulfate and concentrated to yield 2-[(4-bromophenyl)methyl]-cyclohexanone dimethylhydrazone (16 g) as a yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.35 (2H, d), 7.05 (2H, d), 3.05 (1H, dd), 2.4 (6H, s), 2.7-2.2 (4H, m), 1.8-1.3 (6H, m).

### Preparation 4


#### **2-[2-[(4-bromophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium, iodide**



2-[(4-bromophenyl)methyl]-cyclohexanone dimethylhydrazone (16.2 g) (0.052 mol) and methyl iodide (60 ml) were stirred overnight at RT. The precipitate was filtered, washed twice with ethyl ether, then with EtOH, and finally with pentane. After drying, 2-[2-[(4-bromophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium iodide was obtained as a white solid (11.9 g, mp = 173°C). <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.40 (2H, d), 7.05 (2H, d), 3.7 (9H, s), 3.3-3.1 (1H, m), 3.05 (1H, dd), 2.85-2.45 (3H, m), 2.15-1.4 (6H, m).

#### Preparation 5

2-[2-[(4-cyanophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium, iodide




4-[[2-(dimethylhydrazono)cyclohexyl]methyl]-benzonitrile (6.1 g) (0.024 mol) and methyl iodide (35 ml) were stirred overnight at RT. The precipitate was filtered, washed twice with ethyl ether, then with EtOH, and finally with pentane. After drying, 2-[2-[(4-cyanophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium iodide was obtained as a white solid (7.8 g, mp = 175°C). <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 3.7 (9H, s), 3.3 (1H, m), 3.15 (1H, dd), 2.8-2.5 (3H, m), 2.2-1.4 (6H, m).

#### Preparation 6

4-(1-amino-2-oxocyclohexylmethyl)-benzonitrile



47




An EtONa/EtOH solution (from Na [0.42 g][0.0185 mol] in EtOH [15 ml]) was heated to 80°C and 2-[2-[(4-cyanophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium iodide (7.8 g) (0.0185 mol) added in portions. After 1h, the solid has solubilized. The reaction mixture was cooled to RT and a solution of HCl 4M (7.9 ml) carefully added dropwise. After concentration *in vacuo*, the residue was dissolved in EtOAc, extracted twice with HCl 1N, and the aqueous layer washed with tBuOMe. After basification with an NaOH 1N solution, the aqueous phase was extracted with EtOAc. The organic layer washed with water, dried over magnesium sulfate and concentrated to yield 4-[(1-amino-2-oxocyclohexyl)methyl]-benzonitrile (2.46 g) as a brown oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 3.10 (1H, d), 2.85 (1H, d), 2.6 (2H, m), 2.1-1.3 (7H, m).

15

### Preparation 7

#### **2-amino-2-(4-bromo-benzyl)-cyclohexanone**




An EtONa/EtOH solution (from Na [0.57 g][0.025 mol] in EtOH [22 ml]) was heated to 80°C and 2-[2-[(4-bromophenyl)methyl]-cyclohexylidene]-1,1,1-trimethyl-hydrazinium iodide (11.9 g) (0.025 mol) added in portions. After 1h, the solid solubilized. The reaction mixture was cooled to RT and a solution of HCl 4M (10.7 ml) carefully added dropwise. After concentration *in vacuo*, the residue was dissolved in EtOAc, extracted twice with HCl 1N, and the aqueous layer washed with tBuOMe. After basification with an NaOH 1N solution, the aqueous phase was extracted with

EtOAc. The organic layer was washed with water, dried over magnesium sulfate and concentrated to yield 2-amino-2-(4-bromo-benzyl)-cyclohexanone (3.6 g) as a brown oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 7.30 (2H, d), 6.95 (2H, d), 2.98 (1H, d), 2.7 (2H, d), 2.65-2.4 (2H, m), 2.1-1.3 (6H, m +  $\text{NH}_2$ ).

5

### Preparation 8

#### 1-[1-(4-bromo-benzyl)-2-oxocyclohexyl]-3-(3,5-dichlorophenyl)-urea




10

A solution of (3,5-dichlorophenyl)-isocyanate (3.1 g) (1.3 eq) in DCM (30 ml) was carefully dropped on 2-amino-2-(4-bromo-benzyl)-cyclohexanone (3.6 g) (0.013 mol) in DCM (35 ml) at 5°C. The temperature was allowed to warm to 20°C, and the mixture was stirred one night. DCM was evaporated *in vacuo* and the residue purified on  $\text{SiO}_2$  chromatography (eluent DCM/AcOEt - 9.5/0.5) to yield 1-[1-(4-bromo-benzyl)-2-oxocyclohexyl]-3-(3,5-dichlorophenyl)-urea (5.0 g) as a white solid.  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ ): 7.6-7.4 (5H, m), 7.2 (2H, d), 7.05 (1H, bs), 6.6 (1H, bs), 3.0 (1H, d), 2.85 (1H, d), 1.9-1.55 (3H, m), 1.5-1.05 (5H, m).

20

### Preparation 9

#### N-[1-[(4-cyanophenyl)methyl]-2-oxocyclohexyl]-N'-(3,5-dichlorophenyl)-urea




A solution of (3,5-dichlorophenyl)-isocyanate (2.7 g) (1.3 eq) in DCM (25 ml) was carefully dropped on 2-amino-2-(4-cyano-benzyl)-cyclohexanone (2.46 g) (0.011 mol) in DCM (25 ml) at 5°C. The temperature was allowed to warm to 20°C, and the mixture was stirred 6 h. DCM was evaporated *in vacuo* and the residue purified on SiO<sub>2</sub> chromatography (eluent DCM/AcOEt - 9.5/0.5) to yield 1-[1-(4-cyano-benzyl)-2-oxocyclohexyl]-3-(3,5-dichlorophenyl)-urea (3.76 g) as a white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 7.75 (2H, d), 7.45 (2H, d), 7.4 (1H, m), 7.35 (2H, m), 7.1 (1H, bs), 6.6 (1H, bs), 3.1 (1H, d), 2.9 (2H, d), 1.9-1.6 (3H, m), 1.5-1.1 (8H, m).

10

### EXAMPLE 1


#### 3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one



15

A suspension of 1-[1-(4-bromo-benzyl)-2-oxocyclohexyl]-3-(3,5-dichlorophenyl)-urea (5.0 g) (0.0106 mol), K<sub>2</sub>CO<sub>3</sub> (3.1 g) (2.1 eq) and molecular sieves 4A (15 g) in xylene (700 ml) was refluxed for 8 h. After cooling, the reaction mixture was filtered on paper, concentrated *in vacuo*, and the residue purified on SiO<sub>2</sub> chromatography (eluent EtOAc/Cyclohexane - 3/7) to yield 3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (3.2 g) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.45 (2H, d), 7.20 (1H, m), 7.15 (2H, d), 7.05 (2H, m), 4.95 (1H, m), 3.0 (1H, d), 2.90 (1H, d), 2.5-2.1 (2H, m), 2.1-1.8 (3H, m), 1.8-1.6 (1H, m).


25

**EXAMPLE 2****4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile**

5

A suspension of 1-[1-(4-cyano-benzyl)-2-oxocyclohexyl]-3-(3,5-dichlorophenyl)-urea (3.7 g) (0.009 mol), K<sub>2</sub>CO<sub>3</sub> (2.6 g) (2.1 eq) and molecular sieves 4A (10 g) in xylene (500 ml) was refluxed 16 h. After cooling, the reaction mixture was filtered on paper, concentrated *in vacuo*, and the residue purified on SiO<sub>2</sub> chromatography (eluent EtOAc/cyclohexane – 3/7) to yield 4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (1.8 g) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.6 (2H, d), 7.35 (2H, d), 7.15 (1H, m), 6.95 (2H, m), 5.65 (1H, bs), 4.95 (1H, m), 3.1 (1H, d), 2.9 (1H, d), 2.5-2.1 (2H, m), 2.1-1.65 (4H, m).

15


**EXAMPLE 3****4-[1-(3,5-Dichlorophenyl)-3-methyl-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile**

20

4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (200 mg) (0.5 mmol) in DMF (6 ml) was carefully dropped on a suspension of NaH 60% (24.1 mg) (1.2 eq) in DMF (5 ml) at RT. The reaction mixture was stirred 1 h at RT then methyl iodide (0.095 ml) (3 eq) was added. After one night, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to yield a solid which was washed with diisopropyl ether to give 4-[1-(3,5-Dichlorophenyl)-3-methyl-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (115 mg) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 7.10 (1H, m), 6.75 (2H, m), 4.95 (1H, m), 3.10-3.0 (5H, m), 2.45-1.65 (6H, m).

#### EXAMPLE 4


##### 3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-3-methyl-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one



3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (3.0 g) (0.0066 mol) in DMF (50 ml) was carefully dropped on a suspension of NaH 60% (0.32 g) (1.2 eq) in DMF (20 ml) at RT. The reaction mixture was stirred 1 h at RT then methyl iodide (0.6 ml) (1.5 eq) was added. After one night, the reaction mixture was partitioned between water and tBuOMe. The obtained precipitate was filtered, washed with water and tBuOMe, and dried to give 3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-3-methyl-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (1.8 g) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.39 (2H, d), 7.10 (1H, m), 7.0 (2H, d), 6.77 (2H, m), 4.90 (1H, m), 3.0 (3H, s), 2.95 (2H, s), 2.5-2.2 (2H, m), 2.2-1.9 (3H, m), 1.8-1.6 (1H, m).

**EXAMPLE 5****4-[3-Acetyl-1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile**


5



4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (150 mg) (0.38 mmol) in acetic anhydride (5 ml) was refluxed 3  
10 days. After cooling, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was chromatographed over silica gel (DCM/acetone 97.5/2.5), then HPLC to give 4-[3-Acetyl-1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (13.5 mg) as a white solid. <sup>1</sup>H  
15 NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.3-7.2 (3H, m), 6.70 (2H, m), 4.95 (1H, m), 3.70 (1H, d), 3.05 (1H, m), 2.50 (3H, s), 2.45-2.1 (2H, m), 2.1-1.6 (3H, m).

**EXAMPLE 6****3-Acetyl-3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-3-methyl-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one**


20



3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (150 mg) (0.38 mmol) in acetic anhydride (5 ml) was refluxed 3 days. After cooling, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was chromatographed over silica gel (DCM/acetone 97.5/2.5), then HPLC to give 3-Acetyl-3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-3-methyl-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (13.5 mg) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.40 (2H, d), 7.25 (1H, m), 6.98 (2H, d), 6.75 (2H, m), 4.90 (1H, m), 3.55 (1H, d), 3.05-2.9 (2H, m), 2.55 (3H, s), 2.9-2.7 (2H, m), 2.1-1.7 (3H, m).

#### EXAMPLE 7

4-[1-(3,5-Dichlorophenyl)-3-ethyl-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile




4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (150 mg) (0.38 mmol) in DMF (5 ml) was carefully dropped on a suspension of NaH 60% (18.2 mg) (1.2 eq) in DMF (5 ml) at RT. The reaction mixture was stirred 1 h at RT, then ethyl iodide (0.046 ml) (1.5 eq) was added. After one night, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to yield a solid which was purified by HPLC to give 4-[1-(3,5-Dichlorophenyl)-3-ethyl-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (18.2 mg) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 7.10 (1H, m), 6.80 (2H, m), 4.95 (1H, m), 3.75 (1H, m), 3.15 (1H, m), 3.0 (2H, s), 2.5-1.7 (6H, m), 1.25 (3H, t).

**EXAMPLE 8**

**[7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid ethyl ester**

5




4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (150 mg) (0.38 mmol) in DMF (5 ml) was carefully dropped on  
10 a suspension of NaH 60% (18.2 mg) (1.2 eq) in DMF (5 ml) at RT. The reaction mixture was stirred for 1 h at RT, then ethyl bromoacetate (0.052 ml) (1.2 eq) was added. After one night, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified on SiO<sub>2</sub> chromatography (eluent  
15 EtOAc/cyclohexane - 2/8) to give [7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid ethyl ester (46.5 mg) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 7.1 (1H, m), 6.8 (2H, m), 5.0 (1H, m), 4.45 (1H, d), 4.2 (2H, q), 3.75 (1H, d), 3.05 (1H, d), 2.90 (1H, d), 2.5-2.1 (2H, m), 2.1-1.8 (4H, m), 1.25 (3H, t).

20

**EXAMPLE 9**

**[7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid**






[7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid ethyl ester (34 mg) (0.07 mmol) and LiOH (3.0 mg) (1.1 eq) in a 1:1 mixture of MeOH/H<sub>2</sub>O (4 ml) (5 ml) were stirred 4 days. The reaction mixture was acidified with an aqueous solution of HCl 1N, then concentrated. The residue was purified on HPLC to give [7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid (3.9 mg) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 7.15 (1H, m), 6.75 (2H, m), 5.0 (1H, m), 4.45 (1H, d), 3.83 (1H, d), 3.05 (1H, d), 2.9 (1H, d), 2.5-2.1 (2H, m), 2.1-1.8 (4H, m).

#### EXAMPLE 10

[7a-(4-Bromo-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid ethyl ester




3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (190 mg) (0.42 mmol) in DMF (5 ml) was carefully dropped on a suspension of NaH 60% (20.2 mg) (1.2 eq) in DMF (5 ml) at RT. The reaction mixture was stirred 1 h at RT, then ethyl bromoacetate (0.094 ml) (2.0 eq) was added. After one night at RT, the mixture was heated to 100°C for 8 h, then the reaction


mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified on SiO<sub>2</sub> chromatography (eluent EtOAc/cyclohexane – 1/10) to give [7a-(4-Bromo-benzyl)-3-(3,5-dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-acetic acid ethyl ester (30.2 mg) as a yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.35 (2H, d), 7.10 (1H, m), 6.95 (2H, d), 6.8 (2H, m), 4.95 (1H, m), 4.45 (1H, d), 4.2 (2H, q), 3.7 (1H, d), 2.9 (1H, d), 2.8 (1H, d), 2.45-2.1 (2H, m), 2.0-1.8 (4H, m), 1.25 (3H, t).

### EXAMPLE 11

6-[7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-hexanoic acid ethyl ester



4-[1-(3,5-Dichlorophenyl)-2-oxo-1,2,3,4,5,6-hexahydro-benzimidazol-3a-ylmethyl]benzonitrile (500 mg) (1.25 mmol) in DMF (20 ml) was carefully dropped on a suspension of NaH 60% (65 mg) (1.3 eq) in DMF (10 ml) at RT. The reaction mixture was stirred 1 h at RT; then ethyl hexanoate (0.27 ml) (1.2 eq) was added. After 3h, the reaction mixture was poured on water and extracted with tBuOMe. The organic layer was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified on SiO<sub>2</sub> chromatography (eluent EtOAc/cyclohexane - 2/8) to give 6-[7a-(4-Cyano-benzyl)-3-(3,5-Dichlorophenyl)-2-oxo-2,3,5,6,7,7a-hexahydro-benzimidazol-3a-yl]-hexanoic acid ethyl ester (200 mg) as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.55 (2H, d), 7.25 (2H, d), 7.1 (1H, m), 6.75 (2H, m), 4.95 (1H, m), 4.15 (2H, q), 3.7 (1H, m), 3.1 (1H, m), 3.0 (2H, s), 2.35 (2H, m), 2.2-1.3 (12H, m), 1.23 (3H, t).

**EXAMPLE 12****1-(3,5-Dichlorophenyl)-3-methyl-3a-(4-pyrimidin-5-yl-benzyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one**


5

To 3a-(4-Bromo-benzyl)-1-(3,5-dichlorophenyl)-3-methyl-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (1.0 g) (0.002 mol) and 4-pyrimidinyl-trimethylstannate (0.73 g) (1.5 eq) in dry toluene (20 ml) was added tetrakis  
10 (triphenylphosphine) palladium (347 mg) (0.15 eq). The reaction mixture was refluxed 11 h. After cooling, it was filtered on paper and the filtrate concentrated. The residue was purified on SiO<sub>2</sub> chromatography (eluent DCM/acetone – 9/1) then HPLC to give 1-(3,5-Dichlorophenyl)-3-methyl-3a-(4-pyrimidin-5-yl-benzyl)-1,3,3a,4,5,6-hexahydro-benzimidazol-2-one (15 mg) as an off-white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>):  
15 9.15 (1H, s), 8.85 (2H, s), 7.48 (2H, d), 7.26 (2H, d), 7.02 (1H, m), 6.73 (2H, m), 4.93 (1H, m), 3.05 (3H, s), 2.98 (2H, dd), 2.5-1.9 (5H, m), 1.75 (1H, m).

CLAIMS

We claim:

1. A compound of formula (I),



its enantiomers, diastereomers, or a pharmaceutically-acceptable salt, hydrate, or prodrug thereof, in which:

K is O or S;

- Q is a bond,  $-C(=O)-$  or branched or straight chain  $C_{1-4}$ alkylene optionally substituted with one to two  $R_4$ ;

Ar is optionally-substituted aryl or heteroaryl;

$J_1$  is a bond,  $-N(R_5)-$ , or  $-C(R_{6a}R_{7a})-$ ;

$J_2$  is  $-N(R_5)-$  or  $-C(R_{6b}R_{7b})-$ ;

- 15  $J_3$  is  $-N(R_5)-$  or  $-C(R_{6c}R_{7c})-$ ;

provided, however, that only one of  $J_1$ ,  $J_2$  and  $J_3$  may be  $-N(R_5)-$ , so that ring A is a five-to-six membered cycloalkyl or heterocyclo ring having from 0 to 2 heteroatoms;

Y is N or  $C(R_8)$ ;

- 20 Z is N or  $C(R_9)$ ;

$R_1$  is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl,  $-OR_{10}$ ,  $-NR_{10}R_{11}$ ,  $-C(=O)R_{10}$ ,  $-CO_2R_{10}$ ,  $-C(=O)NR_{10}R_{11}$ ,  $-S(O)_pR_{11a}$ ,  $-SO_2NR_{10}R_{11}$ , cycloalkyl, heterocyclo, aryl, and heteroaryl;

R<sub>2</sub> and R<sub>3</sub> are independently selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl, -SR<sub>12</sub>, -OR<sub>12</sub>, -NR<sub>12</sub>R<sub>13</sub>, -CO<sub>2</sub>R<sub>12</sub>, -C(=O)R<sub>12</sub>, -C(=O)NR<sub>12</sub>R<sub>13</sub>, aryl, heterocyclo, cycloalkyl, and heteroaryl;

- 5 R<sub>4</sub> is selected from OH, O(C<sub>1-4</sub>alkyl), halogen, cyano, CF<sub>3</sub>, OCF<sub>3</sub>, NH<sub>2</sub>, NH(C<sub>1-4</sub>alkyl), and N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

R<sub>5</sub> is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cyano, -OR<sub>14</sub>, -NR<sub>14</sub>R<sub>15</sub>, -C(=O)R<sub>14</sub>, -CO<sub>2</sub>R<sub>14</sub>, -C(=O)NR<sub>14</sub>R<sub>15</sub>, -S(O)<sub>p</sub>R<sub>15a</sub>, -SO<sub>2</sub>NR<sub>14</sub>R<sub>15</sub>, aryl, heterocyclo, cycloalkyl, and heteroaryl; or when R<sub>5</sub> is joined  
10 to atom J<sub>1</sub>, J<sub>2</sub> or J<sub>3</sub>, R<sub>5</sub> may be taken together with one of R<sub>6a</sub>, R<sub>6b</sub> or R<sub>6c</sub> attached to an adjacent atom of ring A to form a fused heterocyclo or heteroaryl ring; or when R<sub>5</sub> is joined to atom J<sub>3</sub>, R<sub>5</sub> may be taken together with R<sub>8</sub> to form a fused heterocyclo ring;

R<sub>6a</sub>, R<sub>6b</sub>, R<sub>6c</sub>, R<sub>7a</sub>, R<sub>7b</sub>, R<sub>7c</sub> and R<sub>8</sub> are independently selected from hydrogen, halogen,  
15 alkyl, substituted alkyl, alkenyl, substituted alkenyl, nitro, cyano, -SR<sub>16</sub>, -OR<sub>16</sub>, -NR<sub>16</sub>R<sub>17</sub>, -C(=O)R<sub>16</sub>, -CO<sub>2</sub>R<sub>16</sub>, -C(=O)NR<sub>16</sub>R<sub>17</sub>, -NR<sub>16</sub>C(=O)R<sub>17</sub>, -NR<sub>16</sub>C(=O)OR<sub>17</sub>, -S(O)<sub>q</sub>R<sub>17a</sub>, -NR<sub>16</sub>SO<sub>2</sub>R<sub>17a</sub>, -SO<sub>2</sub>NR<sub>16</sub>R<sub>17</sub>, aryl, heterocyclo, cycloalkyl, and heteroaryl; or R<sub>6a</sub> with R<sub>7a</sub>, or R<sub>6b</sub> with R<sub>7b</sub>, or R<sub>6c</sub> with R<sub>7c</sub> are taken together to form a keto group (=O) or a spiro cycloalkyl or heterocyclo  
20 ring; or R<sub>6b</sub> taken together with either R<sub>6a</sub> or R<sub>6c</sub> may form a fused benzo, cycloalkyl, heterocyclo, or heteroaryl ring; or R<sub>6c</sub> taken together with R<sub>8</sub> may form a fused cycloalkyl or heterocyclo;

R<sub>9</sub> is selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl, -SR<sub>18</sub>, -OR<sub>18</sub>, -NR<sub>18</sub>R<sub>19</sub>, -CO<sub>2</sub>R<sub>18</sub>, -C(=O)R<sub>18</sub>, -  
25 C(=O)NR<sub>18</sub>R<sub>19</sub>, aryl, heterocyclo, cycloalkyl, and heteroaryl;

R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>, and R<sub>19</sub> (i) are selected independently of each other from hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo; or (ii) any two of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>, and R<sub>19</sub> when attached to the same nitrogen atom may be taken together to  
30 form a heteroaryl or heterocyclo ring, with the remainder of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>,

R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>, and R<sub>19</sub> being selected independently from hydrogen, alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo;

R<sub>11a</sub>, R<sub>15a</sub>, and R<sub>17a</sub> are independently selected from alkyl, substituted alkyl, cycloalkyl, aryl, heteroaryl, and heterocyclo;

5 *p* is 1, 2, or 3; and

*q* is 1, 2, or 3.

2. A compound according to claim 1, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein:

K is O;

Q is a -CH<sub>2</sub>-;

Ar is phenyl optionally substituted one to three R<sub>20</sub>;

15 R<sub>1</sub> is selected from hydrogen, C<sub>1-6</sub>alkyl, -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), or C<sub>1-6</sub>alkyl substituted with one to two of hydroxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), and -N(C<sub>1-4</sub>alkyl)<sub>2</sub>;

20 R<sub>2</sub> and R<sub>3</sub> are selected from halogen, (C<sub>1-4</sub>)alkyl, cyano, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, nitro, phenyloxy, benzyloxy, and phenylthio;

R<sub>20</sub> at each occurrence is independently selected from halogen, C<sub>1-4</sub>alkyl, hydroxy, (C<sub>1-4</sub>)alkoxy, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, cyano, nitro, -CO<sub>2</sub>H, -C(=O)H, -CO<sub>2</sub>(C<sub>1-4</sub>)alkyl, -C(=O)(C<sub>1-4</sub>)alkyl, -C(=O)NH(CH<sub>2</sub>)<sub>r</sub>CO<sub>2</sub>H, -C(=O)NH(CH<sub>2</sub>)<sub>r</sub>CO<sub>2</sub>(C<sub>1-4</sub>alkyl), and S(O)<sub>2</sub>(C<sub>1-4</sub>alkyl); or from phenyl, benzyl, 25 phenyloxy, benzyloxy and heteroaryl in turn optionally substituted with one to two of halogen, C<sub>1-4</sub>alkyl, hydroxy, (C<sub>1-4</sub>)alkoxy, halo(C<sub>1-4</sub>)alkyl, halo(C<sub>1-4</sub>)alkoxy, cyano, nitro, -CO<sub>2</sub>H, -C(=O)H, -CO<sub>2</sub>(C<sub>1-4</sub>)alkyl, and/or -C(=O)(C<sub>1-4</sub>)alkyl; or alternatively, two R<sub>20</sub> groups join together with each other to form a fused benzo ring; and

30 *r* is 1, 2, 3, or 4.

3. A compound according to any one of claims 1 to 2, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein

5 J<sub>1</sub> is a bond or -CHR<sub>6a</sub>-;

J<sub>2</sub> is -CHR<sub>6b</sub>-;

J<sub>3</sub> is -CHR<sub>6c</sub>-;

Y is C(R<sub>8</sub>);

R<sub>6a</sub>, R<sub>6b</sub>, R<sub>6c</sub> and R<sub>8</sub> are independently selected from

10 a) hydrogen, halogen, and cyano;

b) -SR<sub>16</sub>, -OR<sub>16</sub>, -NR<sub>16</sub>R<sub>17</sub>, -C(=O)R<sub>16</sub>, -CO<sub>2</sub>R<sub>16</sub>, -C(=O)NR<sub>16</sub>R<sub>17</sub>, -NR<sub>16</sub>C(=O)R<sub>17</sub>, -NR<sub>16</sub>C(=O)OR<sub>17</sub>, -S(O)<sub>q</sub>R<sub>17a</sub>, -NR<sub>16</sub>SO<sub>2</sub>R<sub>17a</sub>, and -SO<sub>2</sub>NR<sub>16</sub>R<sub>17</sub>; and

15 c) C<sub>1-4</sub>alkyl, phenyl, four to seven membered heterocyclo, C<sub>3-7</sub>cycloalkyl, and five to six membered heteroaryl, each of which in turn is optionally substituted with one to two groups selected from R<sub>22</sub>;

R<sub>16</sub> and R<sub>17</sub> are selected independently of each other from hydrogen, C<sub>1-6</sub>alkyl, phenyl, four to seven membered heterocyclo, C<sub>3-7</sub>cycloalkyl, and five to six membered heteroaryl, each of which in turn is optionally substituted with one  
20 to two groups selected from R<sub>23</sub>;

R<sub>17a</sub> is C<sub>1-6</sub>alkyl, phenyl, four to seven membered heterocyclo, C<sub>3-7</sub>cycloalkyl, five to six membered heteroaryl each of which is optionally substituted with one to two groups selected from R<sub>23</sub>; and

25 R<sub>22</sub> and R<sub>23</sub> are at each occurrence selected independently from halogen, cyano, C<sub>1-4</sub>alkyl, hydroxy, trifluoromethyl, trifluoromethoxy, -O(C<sub>1-4</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -C(=O)NH<sub>2</sub>, -C(=O)NH(C<sub>1-4</sub>alkyl), -C(=O)N(C<sub>1-4</sub>alkyl)<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub>alkyl), -N(C<sub>1-4</sub>alkyl)<sub>2</sub>, hydroxy(C<sub>1-4</sub>)alkyl, methoxy(C<sub>1-4</sub>)alkyl, ethoxy(C<sub>1-4</sub>)alkyl, amino(C<sub>1-4</sub>)alkyl, and halo(C<sub>1-4</sub>)alkyl.


30

4. A compound according to any one of claims 1 to 3, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein:

$R_1$  is hydrogen,  $C_{1-6}$ alkyl,  $-C(=O)(C_{1-6}alkyl)$ , or  $C_{1-6}alkyl$  substituted with one of  $-C(=O)H$ ,  $-C(=O)(C_{1-6}alkyl)$ ,  $-CO_2H$ , or  $-CO_2(C_{1-6}alkyl)$ .

5

5. A compound according to any one of claims 1 to 4, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein Q-Ar together form:




wherein

- 10  $R_{20a}$  and  $R_{20b}$  are independently selected from halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4}alkoxy)$ ,  $halo(C_{1-4}alkyl)$ ,  $halo(C_{1-4}alkoxy)$ , cyano, nitro,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4}alkyl)$ ,  $-C(=O)(C_{1-4}alkyl)$ ,  $-C(=O)NH(CH_2)_rCO_2H$ ,  $-C(=O)NH(CH_2)_rCO_2(C_{1-4}alkyl)$ , and  $S(O)_2(C_{1-4}alkyl)$ ; or from phenyl, benzyl, phenyloxy, benzyloxy and heteroaryl in turn optionally substituted with one to
- 15 two of halogen,  $C_{1-4}$ alkyl, hydroxy,  $(C_{1-4}alkoxy)$ ,  $halo(C_{1-4}alkyl)$ ,  $halo(C_{1-4}alkoxy)$ , cyano, nitro,  $-NH_2$ ,  $-NH(C_{1-4}alkyl)$ ,  $-N(C_{1-4}alkyl)_2$ ,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4}alkyl)$ , and/or  $-C(=O)(C_{1-4}alkyl)$ ; or alternatively, two  $R_{20b}$  groups join together with each other or one  $R_{20b}$  joins together with  $R_{20a}$  to form a fused benzo ring;
- 20  $n$  is 0, 1, or 2; and
- $r$  is 1, 2, 3, or 4.

6. A compound according to any one of claims 1 to 5, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein  $J_1$ ,  $J_2$  and  $J_3$  are each
- 25  $-CH_2-$ .



7. A compound according to any one of claims 1 to 6, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein Y is CH.
- 5 8. A compound according to any one of claims 1 to 7, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein R<sub>2</sub> and R<sub>3</sub> are both halogen.
9. A compound according to any one of claims 1 to 8, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, wherein Z is CH.
- 10 10. A compound having the formula (Ia),



15

its enantiomers, diastereomers, or a pharmaceutically-acceptable salt, hydrate, solvate, or prodrug thereof, in which:

Q is -C(=O)- or -(CHR<sub>4a</sub>)<sub>t</sub>-

Ar is aryl or heteroaryl optionally substituted with one to three R<sub>20</sub>;

20 Y is N or C(R<sub>8</sub>);

Z is N or C(R<sub>9</sub>);

R<sub>1</sub> is selected from hydrogen, C<sub>1-6</sub>alkyl, -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), or C<sub>1-6</sub>alkyl substituted with one to two of hydroxy, -O(C<sub>1-6</sub>alkyl), -C(=O)H, -C(=O)(C<sub>1-6</sub>alkyl), -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1-6</sub>alkyl), -C(=O)NH<sub>2</sub>, -

$C(=O)NH_2$ ,  $-C(=O)NH(C_{1-4}alkyl)$ ,  $-C(=O)N(C_{1-4}alkyl)_2$ ,  $-NH_2$ ,  $-NH(C_{1-4}alkyl)$ , and  $-N(C_{1-4}alkyl)_2$ ;

$R_2$  and  $R_3$  are independently selected from halogen,  $(C_{1-4}alkyl)$ , cyano, halo( $C_{1-4}alkyl$ ), halo( $C_{1-4}alkoxy$ ), nitro, phenyloxy, benzyloxy, and phenylthio;

5  $R_{4a}$  is selected from hydrogen, OH,  $O(CH_3)$ ,  $O(CH_2CH_3)$ , halogen, cyano,  $CF_3$ ,  $OCF_3$ ,  $NH_2$ ,  $NH(CH_3)$ , and  $N(CH_3)_2$ ;

$R_6$  and  $R_8$  at each occurrence are independently selected from (a) halogen, nitro, and cyano; or from (b)  $-SR_{16}$ ,  $-OR_{16}$ ,  $-NR_{16}R_{17}$ ,  $-C(=O)R_{16}$ ,  $-CO_2R_{16}$ ,  $-C(=O)NR_{16}R_{17}$ ,  $-NR_{16}C(=O)R_{17}$ ,  $-NR_{16}C(=O)OR_{17}$ ,  $-S(O)_qR_{17a}$ ,  $-NR_{16}SO_2R_{17a}$ ,  
10 and  $-SO_2NR_{16}R_{17}$ ; or from (c) alkyl, alkenyl, aryl, heterocyclo, cycloalkyl, and heteroaryl, in turn optionally substituted with one to two groups selected from  $R_{22}$ ; and/or (d) two  $R_6$  groups taken together form keto ( $=O$ ), with the remainder of the  $R_6$  groups selected from (a), (b), and (c);

$R_9$  is selected from hydrogen, halogen, nitro, cyano, alkyl, substituted alkyl, alkenyl, substituted alkenyl,  $-SR_{18}$ ,  $-OR_{18}$ ,  $-NR_{18}R_{19}$ ,  $-CO_2R_{18}$ ,  $-C(=O)R_{18}$ ,  $-C(=O)NR_{18}R_{19}$ , aryl, heterocyclo, cycloalkyl, and heteroaryl;  
15

$R_{16}$  and  $R_{17}$  are selected independently of each other from hydrogen,  $C_{1-6}alkyl$ , phenyl, four to seven membered heterocyclo,  $C_{3-7}cycloalkyl$ , and five to six membered heteroaryl, each of which in turn is optionally substituted with one to two groups selected from  $R_{23}$ ;  
20

$R_{17a}$  is  $C_{1-6}alkyl$ , phenyl, four to seven membered heterocyclo,  $C_{3-7}cycloalkyl$ , five to six membered heteroaryl each of which is optionally substituted with one to two groups selected from  $R_{23}$ ;

$R_{20}$  at each occurrence is selected from halogen,  $C_{1-6}alkyl$ , hydroxy,  $(C_{1-4}alkoxy)$ ,  $(C_{1-4}alkylthio)$ , cyano, nitro,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4}alkyl)$ ,  $-C(=O)(C_{1-4}alkyl)$ ,  $-C(=O)NH(CH_2)_rCO_2H$ ,  $-C(=O)NH(CH_2)_rCO_2(C_{1-4}alkyl)$ ,  $S(O)_2(C_{1-4}alkyl)$ , phenyl, benzyl, phenyloxy, benzyloxy, five to six membered heteroaryl,  $C_{3-7}cycloalkyl$ , and four to seven membered heterocyclo, wherein each of the alkyl, alkoxy, and cyclic groups in turn are optionally substituted with one to three of  $R_{24}$ ;  
25  
30

$R_{22}$ ,  $R_{23}$  and  $R_{24}$  are at each occurrence selected independently from halogen, cyano, nitro,  $C_{1-6}$ alkyl, hydroxy, trifluoromethyl, trifluoromethoxy,  $-O(C_{1-6}alkyl)$ ,  $-C(=O)H$ ,  $-C(=O)(C_{1-6}alkyl)$ ,  $-CO_2H$ ,  $-CO_2(C_{1-6}alkyl)$ ,  $-C(=O)NH_2$ ,  $-C(=O)NH_2$ ,  $-C(=O)NH(C_{1-4}alkyl)$ ,  $-C(=O)N(C_{1-4}alkyl)_2$ ,  $-NH_2$ ,  $-NH(C_{1-4}alkyl)$ ,  $-N(C_{1-4}alkyl)_2$ , hydroxy( $C_{1-4}$ )alkyl, methoxy( $C_{1-4}$ )alkyl, ethoxy( $C_{1-4}$ )alkyl, amino( $C_{1-4}$ )alkyl, and halo( $C_{1-4}$ )alkyl;


$m$  is 0 or 1;

$p$  and  $q$  are independently 1, 2, or 3;

$r$  and  $s$  are 0, 1, 2, 3 or 4; and

$t$  is 0, 1 or 2.

11. A compound according to claim 10 having the formula,




or a pharmaceutically-acceptable salt, hydrate, solvate, or prodrug thereof, in which:

$R_{20a}$  and  $R_{20b}$  are independently selected from halogen,  $C_{1-4}$ alkyl, hydroxy, ( $C_{1-4}$ )alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4}alkyl)$ ,  $-C(=O)(C_{1-4}alkyl)$ ,  $-C(=O)NH(CH_2)_rCO_2H$ ,  $-C(=O)NH(CH_2)_rCO_2(C_{1-4}alkyl)$ , and  $S(O)_2(C_{1-4}alkyl)$ ; or from phenyl, benzyl, phenyloxy, benzyloxy and heteroaryl in turn optionally substituted with one to two of halogen,  $C_{1-4}$ alkyl, hydroxy, ( $C_{1-4}$ )alkoxy, halo( $C_{1-4}$ )alkyl, halo( $C_{1-4}$ )alkoxy, cyano, nitro,  $-NH_2$ ,  $-NH(C_{1-4}alkyl)$ ,  $-N(C_{1-4}alkyl)_2$ ,  $-CO_2H$ ,  $-C(=O)H$ ,  $-CO_2(C_{1-4}alkyl)$ , and/or  $-C(=O)(C_{1-4}alkyl)$ ; or alternatively, two  $R_{20b}$  groups join together with each other or one  $R_{20b}$  joins together with  $R_{20a}$  to form a fused benzo ring; and

$n$  is 0, 1 or 2.

12. A compound according to any one of claims 10 or 11, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, in which  $R_1$  is  $C_{1-4}$  alkyl.
- 5
13. A compound according to any one of claims 10 to 12, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, in which  $R_2$  and  $R_3$  are both halogen.
- 10 14. A compound according to any one of claims 10 to 13, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, in which  $R_{20a}$  is cyano or halogen.
- 15 15. A compound according to any one of claims 10 or 14, to a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, having the formula,



16. A compound according to any one of claims 10 to 15, or a pharmaceutically-acceptable salt, hydrate, prodrug, or enantiomer thereof, in which Z is CH,  $R_1$  is methyl or ethyl, and  $R_{20a}$  is cyano or halogen.
- 20
17. A pharmaceutical composition comprising at least one compound according to any one of claims 1 to 16, and a pharmaceutically acceptable carrier or diluent.

18. A method of inhibiting an LFA-1/ICAM-associated condition in a mammal comprising administering to the mammal a therapeutically-effective amount of a compound according to any one of claims 1 to 15.
- 5 19. The method of claim 18 in which LFA-1/ICAM-associated condition is selected from acute or chronic graft vs host reactions, acute or chronic transplant rejection, multiple sclerosis, rheumatoid arthritis, psoriatic arthritis, osteoarthritis, osteoporosis, diabetes, cystic fibrosis, inflammatory bowel disease, irritable bowel syndrome, Crohn's disease, ulcerative colitis,
- 10 Alzheimer's disease, shock, ankylosing spondylitis, gastritis, conjunctivitis, pancreatitis, multiple organ injury syndrome, myocardial infarction, atherosclerosis, stroke, reperfusion injury, acute glomerulonephritis, vasculitis, thermal injury, necrotizing enterocolitis, granulocyte transfusion associated syndrome, Sjogren's syndrome, eczema, atopic dermatitis, contact dermatitis,
- 15 urticaria, scleroderma, psoriasis, asthma, pulmonary fibrosis, allergic rhinitis, oxygen toxicity, emphysema, chronic bronchitis, acute respiratory distress syndrome, chronic obstructive pulmonary disease (COPD), hepatitis B, hepatitis C, organ-tissue autoimmune disease, autoimmune thyroiditis, uveitis, systemic lupus erythematosus, Addison's disease, autoimmune polyglandular
- 20 disease, and Grave's disease.